


Journal of Applied Microbiology

Volume 125 / Number 5 / 2018

Edited by A. Gilmour

wileyonlinelibrary.com/journal/jam ISSN 1364-5072



WILEY

sfam
society for
applied microbiology

Published for the Society for Applied
Microbiology by John Wiley & Sons

Journal of Applied Microbiology

Chief Editor

A. Gilmour *Agri-Food and Biosciences Institute, Northern Ireland, UK*
E-mail: cejam@btinternet.com

Reviews Editor

P.J. Collier *University of Abertay, Dundee, UK*

Editors

A.M. Ahmed *Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt*

L. Baillie *Welsh School of Pharmacy, Cardiff University, Cardiff, UK*

J.L. Balcazar *Catalan Institute for Water Research, Girona, Spain*

M.D. Barton *University of South Australia, Adelaide, Australia*

L. Bingle *University of Sunderland, Sunderland, UK*

A. Bosch *Universitat de Barcelona, Spain*

O. Braissant *Department of Urology, University Hospital of Basel, Basel, Switzerland*

J. Brojtsch *Department of Microbiology & Immunology, Albert Einstein College of Medicine, New York City, NY, USA*

P. Calik *Middle East Technical University, Ankara, Turkey*

M. Cao *Department of Chemical and Biomolecular Engineering, University of Illinois at Urbana-Champaign, Urbana, IL, USA*

M.L. Chikindas *Department of Food Science, Rutgers University, New Brunswick, NJ, USA*

C. Cooper *University of Sunderland, Sunderland, UK*

D.A. Cowan *Department of Genetics, University of Pretoria, Pretoria*

S.P. Cummings *School of Applied Sciences, Northumbria University, Newcastle upon Tyne, UK*

M. Dow *BioSciences Institute, University College Cork, Ireland*

N. Fegan *Food Science Australia, Brisbane, Australia*

J.E. Flaherty *Department of Science and Mathematics, Coker College, Hartsville, SC, USA*

C. Gilbert *Laboratoire de Microbiologie et Génétique Moléculaire, Université Claud Bernard Lyon, Lyon, France*

J. Glassey *University of Newcastle Upon Tyne, Newcastle Upon Tyne, UK*

M. Gougouli *Department of Food Science & Technology, School of Agriculture, Aristotle University of Thessaloniki, Greece*

I.R. Grant *Queen's University Belfast, UK*

J. Guard *USDA-ARS, Athens, GA, USA*

S.K. Gupta *Bacterial Epidemiology and Antimicrobial Resistance Research Unit, RRC, Athens, GA, USA*

R.N. Haddadin *School of Pharmacy, The University of Jordan, Amman, Jordan*

K.A. Hammer *School of Pathology and Laboratory Medicine, The University of Western Australia, Perth, Australia*

M. Hernández García *University of Southampton, Southampton, UK*

K.S. Howell *Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville Australia*

C.R. Jackson *USDA-ARS Resistance Research Unit, Athens, GA, USA*

K. Kadlec *Friedrich-Loeffler-Institut, Neustadt-Mariensee, Germany*

M. Koenen *The Netherlands*

S. Kumar *Albany Medical College, Albany, USA*

G. LaPointe *Laval University, Quebec, Canada*

L. Laurens *National Renewable Energy Laboratory, Golden, CO, USA*

J.T. Lisle *USGS, Center for Coastal & Watershed Research, St Petersburg, FL, USA*

J.A. McGarvey *USDA-ARS Plant Mycotoxin Research Unit, Albany, CA, USA*

M. Magnani *Department of Food Engineering, Federal University of Paraíba, João Pessoa, Brazil*

J.-Y. Maillard *Cardiff School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Cardiff, UK*

B. Mayo *IPLA-CSIC, Villaviciosa, Spain*

F. Mozzi *CERELA-CONICET, Tucumán, Argentina*

G.-J.E. Nychas *Department of Food Science & Technology, Laboratory of Food Microbiology & Biotechnology of Foods, Athens, Greece*

F. Pinzari *Council for Agricultural Research and Economics Research Centre for Soil-Plant System [CREA-RPS], Rome, Italy*

K.J. Rashamuse *Industrial Bioinnovation Directorate, Department of Science and Technology, SA*

B.B.A. Raymond *Institut de Pharmacologie et de Biologie Structurale, Centre National de la Recherche Scientifique, Toulouse, France*

A.H. Rickard *School of Public Health, University of Michigan, Ann Arbor, USA*

S. Rossetti *National Research Council of Italy (CNR), Roma, Italy*

C.P. Saint *Water Centre for Water Management and Re-use, University of South Australia, Mawson Lakes, Australia*

A.S. Sant'Ana *Department of Food Science, Faculty of Food Engineering, University of Campinas, Brazil*

P. Setlow *University of Connecticut Health Center, Farmington, CT, USA*

R. Seviour *Biotechnology Research Centre, La Trobe University, Bendigo, Australia*

M.B. Taylor *University of Pretoria, Pretoria, South Africa*

K. Thomas *University of Sunderland, Sunderland, UK*

S.K. Tiwari *Department of Genetics, Maharshi Dayanand University, Rohtak, India*

M. Upton *School of Biomedical & Healthcare Sciences, Plymouth University, Plymouth, UK*

V.P. Valdramidis *University of Malta, Malta*

K. Venema *Beneficial Microbes Consultancy, the Netherlands*

T.M. Wassenaar *Molecular Microbiology and Genomics Consultants, Tannenstrasse, Zotzenhein, Germany*

T.J. Welch *USDA-ARS National Center for Cool and Cold Water Aquaculture, Leetown, WV, USA*

G. Wells *Northwestern University, Evanston, IL, US*

J. Wells *USDA-ARS, Meat Animal Research Center, NE, USA*

R. Zdor *Andrews University, Berrien Springs, MI, USA*

Statistical Adviser

R.A. Armstrong *Aston University, Birmingham, UK*