

Strengthened inequalities for the mean width and the ℓ -norm ^{*†}

Károly J. Böröczky[‡], Ferenc Fodor[§], Daniel Hug

March 5, 2021

Abstract

Barthe proved that the regular simplex maximizes the mean width of convex bodies whose John ellipsoid (maximal volume ellipsoid contained in the body) is the Euclidean unit ball; or equivalently, the regular simplex maximizes the ℓ -norm of convex bodies whose Löwner ellipsoid (minimal volume ellipsoid containing the body) is the Euclidean unit ball. Schmuckenschläger verified the reverse statement; namely, the regular simplex minimizes the mean width of convex bodies whose Löwner ellipsoid is the Euclidean unit ball. Here we prove stronger stability versions of these results. We also consider related stability results for the mean width and the ℓ -norm of the convex hull of the support of centered isotropic measures on the unit sphere.

1 Introduction

In geometric inequalities and extremal problems, Euclidean balls and simplices often are the extremizers. A classical example is the isoperimetric inequality which states that Euclidean balls have smallest surface area among convex bodies (compact convex sets with non-empty interior) of given volume in Euclidean space \mathbb{R}^n , and Euclidean balls are the only minimizers. Another example is the Urysohn inequality which expresses the geometric fact that Euclidean balls minimize the mean width of convex bodies of given volume. To introduce the mean width, let $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ denote the scalar product and Euclidean norm in \mathbb{R}^n , and let B^n be the Euclidean

*Accepted version. Published online: January 13, 2021 in the Journal of the London Mathematical Society, <https://doi.org/10.1112/jlms.12429>.

[†]AMS 2020 subject classification. Primary 52A40; Secondary 52A38, 52B12, 26D15.

Key words and phrases. Mean width, ℓ -norm, simplex, extremal problem, John ellipsoid, Löwner ellipsoid, Brascamp-Lieb inequality, mass transportation, stability result, isotropic measure.

[‡]The author was supported by Hungarian National Research, Development and Innovation Office – NKFIH grants 129630 and 132002.

[§]This research of the author was supported by grant TUDFO/47138-1/2019-ITM of the Ministry for Innovation and Technology, Hungary, and by Hungarian National Research, Development and Innovation Office – NKFIH grants 129630.

unit ball centred at the origin with $\kappa_n = V(B^n) = \pi^{n/2}/\Gamma(1 + n/2)$, where $V(\cdot)$ is the volume (Lebesgue measure) in \mathbb{R}^n . For a convex body K in \mathbb{R}^n , the support function $h_K : \mathbb{R}^n \rightarrow \mathbb{R}$ of K is defined by $h_K(x) = \max_{y \in K} \langle x, y \rangle$ for $x \in \mathbb{R}^n$. Then the mean width of K is given by

$$W(K) = \frac{1}{n\kappa_n} \int_{S^{n-1}} (h_K(u) + h_K(-u)) du,$$

where the integration over the unit sphere S^{n-1} is with respect to the $(n-1)$ -dimensional Hausdorff measure (that coincides with the spherical Lebesgue measure in this case).

A prominent geometric extremal problem for which simplices are extremizers has been discovered and explored much more recently. First, recall that there exists a unique ellipsoid of maximal volume contained in K (which is called the John ellipsoid of K), and a unique ellipsoid of minimal volume containing K (which is called the Löwner ellipsoid of K). It has been shown by Ball [4] that simplices maximize the volume of K given the volume of the John ellipsoid of K , and thus simplices determine the extremal “inner” volume ratio. For the dual problem, Barthe [10] proved that simplices minimize the volume of K given the volume of the Löwner ellipsoid of K , hence simplices determine the extremal “outer” volume ratio (see also Lutwak, Yang, Zhang [53, 55]). In all these cases, equality was characterized by Barthe [10].

In this paper, we consider the mean width and the so called ℓ -norm. To define the latter, for a convex body $K \subset \mathbb{R}^n$ containing the origin in its interior, we set

$$\|x\|_K = \min\{t \geq 0 : x \in tK\}, \quad x \in \mathbb{R}^n.$$

Furthermore, we write γ_n for the standard Gaussian measure in \mathbb{R}^n which has the density function $x \mapsto \sqrt{2\pi}^{-n} e^{-\|x\|^2/2}$, $x \in \mathbb{R}^n$, with respect to Lebesgue measure. Then the ℓ -norm of K is given by

$$\ell(K) = \int_{\mathbb{R}^n} \|x\|_K \gamma_n(dx) = \mathbb{E}\|X\|_K,$$

where X is a Gaussian random vector with distribution γ_n . If the polar body of K is denoted by $K^\circ = \{x \in \mathbb{R}^n : \langle x, y \rangle \leq 1 \ \forall y \in K\}$, then we obtain the relation

$$\ell(K) = \frac{\ell(B^n)}{2} W(K^\circ) \tag{1}$$

with

$$\lim_{n \rightarrow \infty} \frac{\ell(B^n)}{\sqrt{n}} = 1.$$

In addition, the ℓ -norm of K can be expressed in the form (see Barthe [11])

$$\ell(K) = \int_{\mathbb{R}^n} \mathbb{P}(\|X\|_K > t) dt = \int_0^\infty (1 - \gamma_n(tK)) dt. \tag{2}$$

Let Δ_n be a regular simplex inscribed into B^n , and hence Δ_n° is a regular simplex circumscribed around B^n . Theorem 1.1 (i) is due to Barthe [11], and (ii) was proved by Schmuckenschläger [62].

Theorem 1.1 (Barthe '98, Schmuckenschläger '99) *Let K be a convex body in \mathbb{R}^n .*

- (i) *If $B^n \supset K$ is the Löwner ellipsoid of K , then $\ell(K) \leq \ell(\Delta_n)$, and if $B^n \subset K$ is the John ellipsoid of K , then $W(K) \leq W(\Delta_n^\circ)$. Equality holds in either case if and only if K is a regular simplex.*
- (ii) *If $B^n \subset K$ is the John ellipsoid of K , then $\ell(K) \geq \ell(\Delta_n^\circ)$, and if $B^n \supset K$ is the Löwner ellipsoid of K , then $W(K) \geq W(\Delta_n)$. Equality holds in either case if and only if K is a regular simplex.*

It follows from (1) and the duality of Löwner and John ellipsoids that the two statements in (i) are equivalent to each other, and the same is true for (ii).

The classical Urysohn inequality states that $(W(K)/2)^n \geq V(K)/\kappa_n$ with equality exactly when K is a ball. While a reverse form of the Urysohn inequality is still not known in general, we recall that Giannopoulos, Milman, Rudelson [33] proved a reverse Urysohn inequality, for zonoids, and Hug, Schneider [43] established reverse inequalities of other intrinsic and mixed volumes for zonoids and explored applications to stochastic geometry. A related classical open problem in convexity and probability theory is that among all simplices contained in the Euclidean unit ball, the inscribed regular simplex has the maximal mean width (see Litvak [51] for a comprehensive survey on this topic).

Let us discuss the range of $W(K)$ (and hence that of $\ell(K)$ by (1)) in Theorem 1.1. If K is a convex body in \mathbb{R}^n whose Löwner ellipsoid is B^n , then the monotonicity of the mean width and Theorem 1.1 (i) yield

$$W(\Delta_n) \leq W(K) \leq W(B^n) = 2,$$

where, according to Böröczky [19], we have

$$W(\Delta_n) \sim 4\sqrt{\frac{2 \ln n}{n}} \text{ as } n \rightarrow \infty.$$

In addition, if K is a convex body in \mathbb{R}^n whose John ellipsoid is B^n , then

$$2 = W(B^n) \leq W(K) \leq W(\Delta_n^\circ)$$

with $W(\Delta_n^\circ) \sim 4\sqrt{2n \ln n}$.

An important concept in the proof of Theorem 1.1 is the notion of an isotropic measure on the unit sphere. Following Giannopoulos, Papadimitrakis [34] and Lutwak, Yang, Zhang [55], we call a Borel measure μ on the unit sphere S^{n-1} isotropic if

$$\int_{S^{n-1}} u \otimes u \mu(du) = I_n, \tag{3}$$

where I_n is the identity map (or the identity matrix). Condition (3) is equivalent to

$$\langle x, x \rangle = \int_{S^{n-1}} \langle u, x \rangle^2 \mu(du) \text{ for } x \in \mathbb{R}^n.$$

In this case, equating traces of the two sides of (3), we obtain that $\mu(S^{n-1}) = n$. In addition, we say that the isotropic measure μ on S^{n-1} is centered if

$$\int_{S^{n-1}} u \mu(du) = o.$$

We observe that if μ is a centered isotropic measure on S^{n-1} , then **for the cardinality $|\text{supp } \mu|$ of the support of μ it holds that $|\text{supp } \mu| \geq n + 1$** , with equality if and only if μ is concentrated on the vertices of some regular simplex and each vertex has measure $n/(n + 1)$ (see [20, Lemma 10.2] for a quantitative version of this fact).

We recall that isotropic measures on \mathbb{R}^n play a central role in the KLS conjecture by Kannan, Lovász and Simonovits [46] as well as in the analysis of Bourgain’s hyperplane conjecture (slicing problem); see, for instance, Barthe and Cordero-Erausquin [13], Guedon and Milman [42], Klartag [47], Artstein-Avidan, Giannopoulos, Milman [2] and Alonso-Gutiérrez, Bastero [1].

The emergence of isotropic measures on S^{n-1} arises from Ball’s crucial insight that John’s characteristic condition [44, 45] for a convex body to have the unit ball as its John or Löwner ellipsoid (see [3, 4]) can be used to give the Brascamp-Lieb inequality a convenient form which is ideally suited for many geometric applications (see Section 2). John’s characteristic condition (with the proof of the equivalence completed by Ball [6]) states that B^n is the John ellipsoid of a convex body K containing B^n if and only if there exist distinct unit vectors $u_1, \dots, u_k \in \partial K \cap S^{n-1}$ and $c_1, \dots, c_k > 0$ such that

$$\sum_{i=1}^k c_i u_i \otimes u_i = I_n, \tag{4}$$

$$\sum_{i=1}^k c_i u_i = o. \tag{5}$$

In particular, the measure μ on S^{n-1} with support $\{u_1, \dots, u_k\}$ and $\mu(\{u_i\}) = c_i$ for $i = 1, \dots, k$ is isotropic and centered. In addition, B^n is the Löwner ellipsoid of a convex body $K \subset B^n$ if and only if there exist $u_1, \dots, u_k \in \partial K \cap S^{n-1}$ and $c_1, \dots, c_k > 0$ satisfying (4) and (5). According to John [45] (see also Gruber, Schuster [40]), we may assume that $k \leq n(n + 3)/2$ in (4) and (5). It follows from John’s characterization that B^n is the Löwner ellipsoid of a convex body $K \subset B^n$ if and only if B^n is the John ellipsoid of K° .

The finite Borel measures on S^{n-1} which have an isotropic linear image are characterized by Böröczky, Lutwak, Yang and Zhang [21], building on earlier work by Carlen, and Cordero-Erausquin [23], Bennett, Carbery, Christ and Tao [17] and Klartag [48].

We write $\text{conv } X$ to denote the convex hull of a set $X \subset \mathbb{R}^n$. We observe that if μ is a centered isotropic measure on S^{n-1} , then $o \in \text{int } Z_\infty(\mu)$ for

$$Z_\infty(\mu) = \text{conv supp } \mu.$$

For the present purpose, the study of $Z_\infty(\mu)$ can be reduced to discrete measures, as Lemma 10.1 in Böröczky, Hug [20] states that for any centered isotropic measure μ , there exists a discrete

centered isotropic measure μ_0 on S^{n-1} whose support is contained in the support of μ (see Lemma 2.1). It follows that Theorem 1.1 is equivalent to the following statements about isotropic measures proved by Li, Leng [49].

Theorem 1.2 (Li, Leng '12) *If μ is a centered isotropic measure on S^{n-1} , then $\ell(Z_\infty(\mu)) \leq \ell(\Delta_n)$, $W(Z_\infty(\mu)^\circ) \leq W(\Delta_n^\circ)$, $\ell(Z_\infty(\mu)^\circ) \geq \ell(\Delta_n^\circ)$ and $W(Z_\infty(\mu)) \geq W(\Delta_n)$, with equality in either case if and only if $|\text{supp } \mu| = n + 1$.*

Results similar to Theorem 1.2 are proved by Ma [56] in the L_p setting.

The main goal of the present paper is to provide stronger stability versions of Theorem 1.1 and Theorem 1.2. Since our results use the notion of distance between convex bodies (and to fix the notation), we recall that the distance between compact subsets X and Y of \mathbb{R}^n is measured in terms of the Hausdorff distance defined by

$$\delta_H(X, Y) = \max\left\{\max_{y \in Y} d(y, X), \max_{x \in X} d(x, Y)\right\},$$

where $d(x, Y) = \min\{\|x - y\| : y \in Y\}$. The Hausdorff distance defines a metric on the set of non-empty compact subsets of \mathbb{R}^n .

In addition, for convex bodies K and C , the symmetric difference distance of K and C is the volume of their symmetric difference; namely,

$$\delta_{\text{vol}}(K, C) = V(K \setminus C) + V(C \setminus K).$$

Clearly, the symmetric difference distance also defines a metric on the set of convex bodies in \mathbb{R}^n . Both metrics induce the same topology on the space of convex bodies, **but are not uniformly equivalent to each other (see [63, p. 71] and [64]).**

Let $O(n)$ denote the orthogonal group (rotation group) of \mathbb{R}^n .

Theorem 1.3 *Let B^n be the Löwner ellipsoid of a convex body $K \subset B^n$ in \mathbb{R}^n , let $c = n^{26n}$ and let $\varepsilon \in (0, 1)$. If $\ell(K) \geq (1 - \varepsilon)\ell(\Delta_n)$, then there exists a $T \in O(n)$ such that*

$$(i) \quad \delta_{\text{vol}}(K, T\Delta_n) \leq c \sqrt[4]{\varepsilon},$$

$$(ii) \quad \delta_H(K, T\Delta_n) \leq c \sqrt[4]{\varepsilon}.$$

Theorem 1.4 *Let B^n be the John ellipsoid of a convex body $K \supset B^n$ in \mathbb{R}^n and let $\varepsilon > 0$. If $\ell(K) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$, then there exists a $T \in O(n)$ such that*

$$(i) \quad \delta_{\text{vol}}(K, T\Delta_n^\circ) \leq c \sqrt[4]{\varepsilon} \text{ for } c = n^{27n},$$

$$(ii) \quad \delta_H(K, T\Delta_n^\circ) \leq c \sqrt[4n]{\varepsilon} \text{ for } c = n^{27}.$$

Let us consider the optimality of the order of the estimates in Theorems 1.3 and 1.4. For Theorem 1.3 (i) and (ii) **we use the following construction. We add an $(n + 2)$ nd vertex $v_{n+2} \in S^{n-1}$ to the $n + 1$ vertices v_1, \dots, v_{n+1} of Δ_n such that v_1 lies on the geodesic arc on S^{n-1} connecting v_2 and v_{n+2} , and such that $\angle(v_{n+2}, v_1) = c_1\varepsilon$ for a suitable $c_1 > 0$ depending on**

n . The polytope $K = \text{conv}\{v_1, \dots, v_{n+2}\}$ satisfies $\ell(K) \geq (1 - \varepsilon)\ell(\Delta_n)$ on the one hand, and $\delta_{\text{vol}}(K, T\Delta_n) \geq c_2\varepsilon$ and $\delta_H(K, T\Delta_n) \geq c_2\varepsilon$ for a suitable $c_2 > 0$, depending on n , and for any $T \in O(n)$, on the other hand. Similarly, using the polar of this polytope K for Theorem 1.4 (i), possibly after decreasing c_1 , we have $\ell(K^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ while $\delta_{\text{vol}}(K^\circ, T\Delta_n^\circ) \geq c_3\varepsilon$ for a suitable $c_3 > 0$ depending on n and for any $T \in O(n)$. Finally, we consider the optimality of Theorem 1.4 (ii). Cutting off $n + 1$ regular simplices of edge length $c_4 \sqrt[n]{\varepsilon}$ at the vertices of Δ_n° , for a suitable $c_4 > 0$ depending on n , results in a polytope \tilde{K} satisfying $\ell(\tilde{K}) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ and $\delta_H(\tilde{K}, T\Delta_n^\circ) \geq c_5 \sqrt[n]{\varepsilon}$ for any $T \in O(n)$ for some suitable $c_5 > 0$ depending on n .

We did not make an attempt to optimize the constants c that depend on n , but observe that the c is polynomial in n in Theorem 1.4 (ii).

In the case of the mean width, we have the following stability versions of Theorem 1.1.

Corollary 1.5 *Let K be convex body in \mathbb{R}^n .*

- (i) *If B^n is the John ellipsoid of $K \supset B^n$ and $W(K) \geq (1 - \varepsilon)W(\Delta_n^\circ)$ for some $\varepsilon \in (0, 1)$, then there exists a $T \in O(n)$ such that $\delta_H(K, T\Delta_n^\circ) \leq c\sqrt[n]{\varepsilon}$ for $c = n^{27n}$.*
- (ii) *If B^n is the Löwner ellipsoid of $K \subset B^n$ and $W(K) \leq (1 + \varepsilon)W(\Delta_n)$ for some $\varepsilon > 0$, then there exists a $T \in O(n)$ such that $\delta_H(K, T\Delta_n) \leq c\sqrt[n]{\varepsilon}$ for $c = n^{29}$.*

For the optimality of Corollary 1.5 (i), cutting off $n + 1$ regular simplices of edge length $c_1\varepsilon$ at the vertices of Δ_n° for suitable $c_1 > 0$ depending on n results in a polytope K satisfying $W(K) \geq (1 - \varepsilon)W(\Delta_n^\circ)$ and $\delta_H(K, T\Delta_n^\circ) \geq c_2\varepsilon$ for suitable $c_2 > 0$ depending on n and for any $T \in O(n)$. Concerning Corollary 1.5 (ii), let v_1, \dots, v_{n+1} be the vertices of Δ_n , and let \tilde{K} be the polytope whose vertices are $v_i, -(\frac{1}{n} + c_3 \sqrt[n]{\varepsilon})v_i$ for $i = 1, \dots, n + 1$ for suitable $c_3 > 0$ depending on n in a way such that $W(\tilde{K}) \leq (1 + \varepsilon)W(\Delta)$. It follows that $\delta_H(K, T\Delta_n) \geq c_4 \sqrt[n]{\varepsilon}$ for any $T \in O(n)$ and for a suitable $c_4 > 0$ depending on n .

We also have the following stronger form of Theorem 1.2 in the form of stability statements.

Theorem 1.6 *Let μ be a centered isotropic measure on the unit sphere S^{n-1} , let $c = n^{28n}$, and let $\varepsilon \in (0, 1)$. If one of the conditions*

- (a) $\ell(Z_\infty(\mu)) \geq (1 - \varepsilon)\ell(\Delta_n)$ or
- (b) $W(Z_\infty(\mu)^\circ) \geq (1 - \varepsilon)W(\Delta_n^\circ)$ or
- (c) $\ell(Z_\infty(\mu)^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ or
- (d) $W(Z_\infty(\mu)) \leq (1 + \varepsilon)W(\Delta_n)$

is satisfied, then there exists a regular simplex with vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ such that

$$\delta_H(\text{supp } \mu, \{w_1, \dots, w_{n+1}\}) \leq c\varepsilon^{\frac{1}{4}}.$$

The proofs of Theorem 1.3 and Theorem 1.6 (a), (b) are based on Proposition 7.1, which is the special case of Theorem 1.6 (a) for a discrete measure. In addition, a new stability version of Barthe's reverse of the Brascamp-Lieb inequality is required for a special parametric class of functions, which is derived in Section 6. In a similar vein, the proofs of Theorem 1.4 and Theorem 1.6 (c), (d) are based on Proposition 9.1, which is the special case of Theorem 1.6 (c) for a discrete measure. In addition, we use and derive a stability version of the Brascamp-Lieb inequality for a special parametric class of functions (see also Section 6).

We note that our arguments are based on the rank one geometric Brascamp-Lieb and reverse Brascamp-Lieb inequalities (see Section 2), and their stability versions in a special case (see Section 6). Unfortunately, no quantitative stability version of the Brascamp-Lieb and reverse Brascamp-Lieb inequalities are known in general (see Bennett, Bez, Flock, Lee [16] for a certain weak stability version for higher ranks). On the other hand, in the case of the Borell-Brascamp-Lieb inequality (see Borell [18], Brascamp, Lieb [22] and Balogh, Kristály [8]), stability versions were proved by Ghilli, Salani [32] and Rossi, Salani [60].

2 Discrete isotropic measures and the (reverse) Brascamp-Lieb inequality

For the purposes of this paper, the study of $Z_\infty(\mu)$ for centered isotropic measures on S^{n-1} can be reduced to the case when μ is discrete. Writing $|X|$ for the cardinality of a finite set X , we recall that Lemma 10.1 in Böröczky, Hug [20] states that for any centered isotropic measure μ , there exists a discrete centered isotropic measure μ_0 on S^{n-1} with $\text{supp } \mu_0 \subset \text{supp } \mu$ and $|\text{supp } \mu_0| \leq \frac{n(n+3)}{2} + 1$. We use this statement in the following form.

Lemma 2.1 *For any centered isotropic measure μ on S^{n-1} , there exists a discrete centered isotropic measure μ_0 on S^{n-1} such that*

$$\text{supp } \mu_0 \subset \text{supp } \mu \text{ and } |\text{supp } \mu_0| \leq 2n^2.$$

The rank one geometric Brascamp-Lieb inequality (7) was identified by Ball [3] as an important case of the rank one Brascamp-Lieb inequality proved originally by Brascamp, Lieb [22]. In addition, the reverse Brascamp-Lieb inequality (8) is due to Barthe [9, 10]. To set up (7) and (8), let the distinct unit vectors $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ satisfy

$$\sum_{i=1}^k c_i u_i \otimes u_i = I_n. \quad (6)$$

If f_1, \dots, f_k are non-negative measurable functions on \mathbb{R} , then the Brascamp-Lieb inequality states that

$$\int_{\mathbb{R}^n} \prod_{i=1}^k f_i(\langle x, u_i \rangle)^{c_i} dx \leq \prod_{i=1}^k \left(\int_{\mathbb{R}} f_i \right)^{c_i}, \quad (7)$$

and the reverse Brascamp-Lieb inequality is given by

$$\int_{\mathbb{R}^n}^* \sup_{x=\sum_{i=1}^k c_i \theta_i u_i} \prod_{i=1}^k f_i(\theta_i)^{c_i} dx \geq \prod_{i=1}^k \left(\int_{\mathbb{R}} f_i \right)^{c_i}, \quad (8)$$

where the star on the left-hand-side denotes the upper integral. Here we always assume that $\theta_1, \dots, \theta_k \in \mathbb{R}$ in (8). We note that $\theta_1, \dots, \theta_k$ are unique if $k = n$ and hence u_1, \dots, u_n is an orthonormal basis.

It was proved by Barthe [10] that equality in (7) or in (8) implies that if none of the functions f_i is identically zero or a scaled version of a Gaussian, then there exists an origin symmetric regular crosspolytope in \mathbb{R}^n such that u_1, \dots, u_k lie among its vertices. Conversely, we note that equality holds in (7) and (8) if either each f_i is a scaled version of the same centered Gaussian, or if $k = n$ and u_1, \dots, u_n form an orthonormal basis.

For a detailed discussion of the rank one Brascamp-Lieb inequality, we refer to Carlen, Cordero-Erausquin [23]. The higher rank case, due to Lieb [50], is reproved and further explored by Barthe [10]. Equality in the general version of the Brascamp-Lieb inequality is clarified by Bennett, Carbery, Christ, Tao [17]. In addition, Barthe, Cordero-Erausquin, Ledoux, Maurey (see [14]) develop an approach for the Brascamp-Lieb inequality via Markov semigroups in a quite general framework.

The fundamental papers by Barthe [9, 10] provided concise proofs of (7) and (8) based on mass transportation (see Ball [7] for a sketch in the case of (7)). Actually, the reverse Brascamp-Lieb inequality (8) seems to be the first inequality whose original proof is via mass transportation. During the argument in Barthe [10], the following four observations due to K.M. Ball [3] (see also [10] for a simpler proof of (i)) play crucial roles: If $k \geq n$, $c_1, \dots, c_k > 0$ and $u_1, \dots, u_k \in S^{n-1}$ satisfy (6), then

(i) for any $t_1, \dots, t_k > 0$, we have

$$\det \left(\sum_{i=1}^k t_i c_i u_i \otimes u_i \right) \geq \prod_{i=1}^k t_i^{c_i}, \quad (9)$$

(ii) for $z = \sum_{i=1}^k c_i \theta_i u_i$ with $\theta_1, \dots, \theta_k \in \mathbb{R}$, we have

$$\|z\|^2 \leq \sum_{i=1}^k c_i \theta_i^2, \quad (10)$$

(iii) for $i = 1, \dots, k$, we have

$$c_i \leq 1,$$

(iv) and it holds that

$$c_1 + \dots + c_k = n. \quad (11)$$

Inequality (9) is called the Ball-Barthe inequality by Lutwak, Yang, Zhang [55] and Li, Leng [49].

3 Review of the proof of the (reverse) Brascamp-Lieb inequality if all $f_i = f$ and f is log-concave

Let $g(t) = \sqrt{2\pi}^{-1} e^{-t^2/2}$, $t \in \mathbb{R}$, be the standard Gaussian density (mean zero, variance one), and let f be a probability density function on \mathbb{R} (here we restrict to log-concave functions to avoid differentiability issues). Let T and S be the transportation maps which are determined by

$$\int_{-\infty}^x f = \int_{-\infty}^{T(x)} g \quad \text{and} \quad \int_{-\infty}^{S(y)} f = \int_{-\infty}^y g.$$

Henceforth, we do not write the arguments and the Lebesgue measure in the integral if the meaning of the integral is unambiguous. As f is log-concave, there exists an open interval I such that f is positive on I and zero on the complement of the closure of I , and $T : I \rightarrow \mathbb{R}$ and $S : \mathbb{R} \rightarrow I$ are inverses of each other. In addition, for $x \in I$ and $y \in \mathbb{R}$ we have

$$f(x) = g(T(x)) T'(x) \quad \text{and} \quad g(y) = f(S(y)) S'(y). \quad (12)$$

For

$$\mathcal{C} = \{x \in \mathbb{R}^n : \langle u_i, x \rangle \in I \text{ for } i = 1, \dots, k\},$$

we consider the transformation $\Theta : \mathcal{C} \rightarrow \mathbb{R}^n$ with

$$\Theta(x) = \sum_{i=1}^k c_i T(\langle u_i, x \rangle) u_i, \quad x \in \mathcal{C},$$

which satisfies

$$d\Theta(x) = \sum_{i=1}^k c_i T'(\langle u_i, x \rangle) u_i \otimes u_i.$$

It is known that $d\Theta$ is positive definite and $\Theta : \mathcal{C} \rightarrow \mathbb{R}^n$ is injective (see [9, 10]). Therefore, using first (12), then (i) with $t_i = T'(\langle u_i, x \rangle)$, and then the definition of Θ and (ii), the following argument leads to the Brascamp-Lieb inequality in this special case:

$$\begin{aligned} & \int_{\mathbb{R}^n} \prod_{i=1}^k f(\langle u_i, x \rangle)^{c_i} dx \\ &= \int_{\mathcal{C}} \prod_{i=1}^k f(\langle u_i, x \rangle)^{c_i} dx \\ &= \int_{\mathcal{C}} \left(\prod_{i=1}^k g(T(\langle u_i, x \rangle))^{c_i} \right) \left(\prod_{i=1}^k T'(\langle u_i, x \rangle)^{c_i} \right) dx \\ &\leq \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathcal{C}} \left(\prod_{i=1}^k e^{-c_i T(\langle u_i, x \rangle)^2/2} \right) \det \left(\sum_{i=1}^k c_i T'(\langle u_i, x \rangle) u_i \otimes u_i \right) dx \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathcal{C}} e^{-\|\Theta(x)\|^2/2} \det(d\Theta(x)) dx \\
&\leq \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^n} e^{-\|y\|^2/2} dy = 1.
\end{aligned}$$

We note that the Brascamp-Lieb inequality (13) for an arbitrary non-negative log-concave function f follows by scaling; namely, (iv) implies

$$\int_{\mathbb{R}^n} \prod_{i=1}^k f(\langle x, u_i \rangle)^{c_i} dx \leq \left(\int_{\mathbb{R}} f \right)^n. \quad (13)$$

For the reverse Brascamp-Lieb inequality, we observe that

$$d\Psi(x) = \sum_{i=1}^k c_i S'(\langle u_i, x \rangle) u_i \otimes u_i$$

holds for the differentiable map $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with

$$\Psi(x) = \sum_{i=1}^k c_i S(\langle u_i, x \rangle) u_i, \quad x \in \mathbb{R}^n.$$

In particular, $d\Psi$ is positive definite and $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is injective (see [9, 10]). Therefore (i) and (12) lead to **(for the first inequality, observe that Ψ is injective, but Ψ need not be surjective)**

$$\begin{aligned}
&\int_{\mathbb{R}^n}^* \sup_{x=\sum_{i=1}^k c_i \theta_i u_i} \prod_{i=1}^k f(\theta_i)^{c_i} dx \\
&\geq \int_{\mathbb{R}^n}^* \left(\sup_{\Psi(y)=\sum_{i=1}^k c_i \theta_i u_i} \prod_{i=1}^k f(\theta_i)^{c_i} \right) \det(d\Psi(y)) dy \\
&\geq \int_{\mathbb{R}^n} \left(\prod_{i=1}^k f(S(\langle u_i, y \rangle))^{c_i} \right) \det \left(\sum_{i=1}^k c_i S'(\langle u_i, y \rangle) u_i \otimes u_i \right) dy \\
&\geq \int_{\mathbb{R}^n} \left(\prod_{i=1}^k f(S(\langle u_i, y \rangle))^{c_i} \right) \left(\prod_{i=1}^k S'(\langle u_i, y \rangle)^{c_i} \right) dy \\
&= \int_{\mathbb{R}^n} \left(\prod_{i=1}^k g(\langle u_i, y \rangle)^{c_i} \right) dy = \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^n} e^{-\|y\|^2/2} dy = 1.
\end{aligned}$$

Again, the reverse Brascamp-Lieb inequality (14) for an arbitrary non-negative log-concave function f follows by scaling and (iv); namely,

$$\int_{\mathbb{R}^n}^* \sup_{x=\sum_{i=1}^k c_i \theta_i u_i} \prod_{i=1}^k f(\theta_i)^{c_i} dx \geq \left(\int_{\mathbb{R}} f \right)^n. \quad (14)$$

We observe that (i) shows that the optimal factor in the geometric Brascamp-Lieb inequality and in its reverse form is 1.

4 Observations on the stability of the Brascamp-Lieb inequality and its reverse

This section **summarizes** certain stability forms of the Ball-Barthe inequality (9) based on work in Böröczky, Hug [20]. The first step is a stability version of the Ball-Barthe inequality (9) proved in [20].

Lemma 4.1 *If $k \geq n + 1$, $t_1, \dots, t_k > 0$, $c_1, \dots, c_k > 0$ and $u_1, \dots, u_k \in S^{n-1}$ satisfy (6), then*

$$\det \left(\sum_{i=1}^k t_i c_i u_i \otimes u_i \right) \geq \theta^* \prod_{i=1}^k t_i^{c_i},$$

where

$$\theta^* = 1 + \frac{1}{2} \sum_{1 \leq i_1 < \dots < i_n \leq k} c_{i_1} \cdots c_{i_n} \det[u_{i_1}, \dots, u_{i_n}]^2 \left(\frac{\sqrt{t_{i_1} \cdots t_{i_n}}}{t_0} - 1 \right)^2,$$

$$t_0 = \sqrt{\sum_{1 \leq i_1 < \dots < i_n \leq k} t_{i_1} \cdots t_{i_n} c_{i_1} \cdots c_{i_n} \det[u_{i_1}, \dots, u_{i_n}]^2}.$$

In order to estimate θ^* from below, we use the following observation from [20].

Lemma 4.2 *If $a, b, x > 0$, then*

$$(xa - 1)^2 + (xb - 1)^2 \geq \frac{(a^2 - b^2)^2}{2(a^2 + b^2)^2}.$$

The combination of Lemma 4.1 and Lemma 4.2 implies the following stability version of the Ball-Barthe inequality (9) that is easier to use.

Corollary 4.3 *If $k \geq n + 1$, $t_1, \dots, t_k > 0$, $c_1, \dots, c_k > 0$ and $u_1, \dots, u_k \in S^{n-1}$ satisfy (6), and there exist $\beta > 0$ and $n + 1$ indices $\{i_1, \dots, i_{n+1}\} \subset \{1, \dots, k\}$ such that*

$$c_{i_1} \cdots c_{i_n} \det[u_{i_1}, \dots, u_{i_n}]^2 \geq \beta,$$

$$c_{i_2} \cdots c_{i_{n+1}} \det[u_{i_2}, \dots, u_{i_{n+1}}]^2 \geq \beta,$$

then

$$\det \left(\sum_{i=1}^k t_i c_i u_i \otimes u_i \right) \geq \left(1 + \frac{\beta(t_{i_1} - t_{i_{n+1}})^2}{4(t_{i_1} + t_{i_{n+1}})^2} \right) \prod_{i=1}^k t_i^{c_i}.$$

We may assume that $k \leq 2n^2$ (see Lemma 2.1), and thus the following observation from [20] can be used to estimate β in Corollary 4.3 from below.

Lemma 4.4 *If $k \geq n$, $c_1, \dots, c_k > 0$ and $u_1, \dots, u_k \in S^{n-1}$ satisfy (6), then there exist $1 \leq i_1 < \dots < i_n \leq k$ such that*

$$c_{i_1} \cdots c_{i_n} \det[u_{i_1}, \dots, u_{i_n}]^2 \geq \binom{k}{n}^{-1}.$$

5 Discrete isotropic measures, orthonormal bases and approximation by a regular simplex

According to Lemma 3.2 and Lemma 5.1 in Böröczky, Hug [20], the following auxiliary results are available.

Lemma 5.1 *Let $v_1, \dots, v_k \in \mathbb{R}^n \setminus \{0\}$ satisfy $\sum_{i=1}^k v_i \otimes v_i = I_n$, and let $0 < \eta < 1/(3\sqrt{k})$. Assume for any $i \in \{1, \dots, k\}$ that $\|v_i\| \leq \eta$ or there is some $j \in \{1, \dots, n\}$ with $\angle(v_i, v_j) \leq \eta$. Then there exists an orthonormal basis w_1, \dots, w_n such that $\angle(v_i, w_i) < 3\sqrt{k}\eta$ for $i = 1, \dots, n$.*

Lemma 5.2 *Let $e \in S^{n-1}$, and let $\tau \in (0, 1/(2n))$. If w_1, \dots, w_n is an orthonormal basis of \mathbb{R}^n such that*

$$\frac{1}{\sqrt{n}} - \tau < \langle e, w_i \rangle < \frac{1}{\sqrt{n}} + \tau \quad \text{for } i = 1, \dots, n,$$

then there exists an orthonormal basis $\tilde{w}_1, \dots, \tilde{w}_n$ such that $\langle e, \tilde{w}_i \rangle = \frac{1}{\sqrt{n}}$ and $\angle(w_i, \tilde{w}_i) < n\tau$ for $i = 1, \dots, n$.

Since $\sqrt{k}(n+1) < kn$ if $k > n \geq 2$, and $|\cos(\beta) - \frac{1}{\sqrt{n+1}}| \leq |\beta - \alpha|$ if $\alpha = \arccos \frac{1}{\sqrt{n+1}}$, we deduce from Lemmas 5.1 and 5.2 the following consequence.

Corollary 5.3 *Let $k > n \geq 2$, let $\tilde{u}_1, \dots, \tilde{u}_k, e \in S^n$ in \mathbb{R}^{n+1} and $\tilde{c}_1, \dots, \tilde{c}_k > 0$ satisfy $\sum_{i=1}^k \tilde{c}_i \tilde{u}_i \otimes \tilde{u}_i = I_{n+1}$ and $\langle e, \tilde{u}_i \rangle = \frac{1}{\sqrt{n+1}}$ for $i = 1, \dots, k$, and let $0 < \eta < 1/(6kn)$. Assume for any $i \in \{1, \dots, k\}$ that $\tilde{c}_i \leq \eta^2$ or there exists some $j \in \{1, \dots, n+1\}$ with $\angle(\tilde{u}_i, \tilde{u}_j) \leq \eta$. Then there exists an orthonormal basis $\tilde{w}_1, \dots, \tilde{w}_{n+1}$ of \mathbb{R}^{n+1} such that $\langle e, \tilde{w}_i \rangle = \frac{1}{\sqrt{n+1}}$ and $\angle(\tilde{u}_i, \tilde{w}_i) < 3kn\eta$ for $i = 1, \dots, n+1$.*

For $\tilde{w}_1, \dots, \tilde{w}_{n+1} \in S^n$ with $\langle e, \tilde{w}_i \rangle = \frac{1}{\sqrt{n+1}}$ for $i = 1, \dots, n+1$, the vectors $\tilde{w}_1, \dots, \tilde{w}_{n+1}$ form an orthonormal basis of \mathbb{R}^{n+1} if and only if their projection to e^\perp form the vertices of a regular n -simplex. Therefore Corollary 5.3 provides information on how close $\text{conv}\{\tilde{u}_1, \dots, \tilde{u}_{n+1}\}$ is to some regular n -simplex. Lemma 5.2 in Böröczky, Hug [20] formulated this observation as follows.

Lemma 5.4 *Let Z be a polytope, and let S be a regular simplex circumscribed to B^n . Assume that the facets of Z and S touch B^n at u_1, \dots, u_k and w_1, \dots, w_{n+1} , respectively. Fix $\eta \in (0, 1/(2n))$. If for any $i \in \{1, \dots, k\}$ there exists some $j \in \{1, \dots, n+1\}$ such that $\angle(u_i, w_j) \leq \eta$, then*

$$(1 - n\eta)S \subset Z \subset (1 + 2n\eta)S.$$

Finally, we need to estimate the difference of Gaussian measures of certain polytopes $Z \subset S$. Since in our case, $S \subset nB^n$, it is equivalent to estimate the volume difference up to a factor depending on n . Our first estimate of this kind is Lemma 5.3 in Böröczky, Hug [20].

Lemma 5.5 *Let Z be a polytope, and let S be a regular simplex both circumscribed to B^n . Fix $\alpha = 9 \cdot 2^{n+2}n^{2n+2}$ and $\eta \in (0, \alpha^{-1})$. Assume that the facets of Z and S touch B^n at u_1, \dots, u_k , $k \geq n+1$, and w_1, \dots, w_{n+1} , respectively. If $\angle(u_i, w_i) \leq \eta$ for $i = 1, \dots, n+1$ and $\angle(u_k, w_i) \geq \alpha\eta$ for $i = 1, \dots, n+1$, then*

$$V(Z) \leq \left(1 - \frac{\min_{i=1, \dots, n+1} \angle(u_k, w_i)}{2^{n+2}n^{2n}}\right) V(S).$$

Secondly, we prove another estimate concerning the volume difference of a convex body and a simplex.

Lemma 5.6 *Let S be a regular simplex whose centroid is the origin, and let $M_1 \subset S$ and $M_2 \supset S$ be convex bodies. Suppose that there is some $\varepsilon \in (0, 1)$ such that $M_1 \not\supset (1 - \varepsilon)S$ for (i) and $(1 + \varepsilon)S \not\subset M_2$ for (ii), respectively. Then*

$$(i) \quad V(S \setminus M_1) \geq \frac{n^n}{(n+1)^n} \varepsilon^n V(S) > \frac{1}{e} \varepsilon^n V(S);$$

$$(ii) \quad V(M_2 \setminus S) \geq \frac{1}{n+1} \varepsilon V(S).$$

Proof. Let R be the circumradius of S , let v_1, \dots, v_{n+1} be the vertices of S , and let u_1, \dots, u_{n+1} be the corresponding exterior unit normals of the facets, and hence

$$v_i = -R u_i \text{ for } i = 1, \dots, n+1, \text{ and } S = \left\{x \in \mathbb{R}^n : \langle x, u_i \rangle \leq \frac{R}{n} \text{ for } i = 1, \dots, n+1\right\}.$$

For (i), there exists a v_i such that $(1 - \varepsilon)v_i \notin M_1$, and hence there exists a closed halfspace H^+ with $(1 - \varepsilon)v_i \in H^+$ and $H^+ \cap M_1 = \emptyset$. We observe that $(1 - \varepsilon)v_i$ is the centroid of the simplex $S_\varepsilon = (1 - \varepsilon)v_i + \varepsilon S \subset S$. Using Grünbaum's result [41] on minimal hyperplane sections of the simplex through its centroid, we obtain

$$V(S \setminus M_1) > V(S_\varepsilon \cap H^+) \geq \frac{n^n}{(n+1)^n} V(S_\varepsilon) = \frac{n^n}{(n+1)^n} \varepsilon^n V(S).$$

For (ii), there exists an $x_0 \in M_2 \setminus ((1 + \varepsilon)S)$, and hence there is a u_j such that $\langle x_0, u_j \rangle > \frac{(1 + \varepsilon)R}{n}$. We write F_j to denote the facet of S with exterior unit normal u_j , and $|F_j|$ to denote the $(n - 1)$ -volume of F_j . It follows that

$$V(M_2 \setminus S) \geq \frac{1}{n} \frac{\varepsilon R}{n} |F_j| = \frac{\varepsilon}{n+1} (n+1) \frac{1}{n} \frac{R}{n} |F_j| = \frac{\varepsilon}{n+1} V(S),$$

which completes the proof. \square

Remark The estimates in (i) and in (ii) are optimal in the sense that there exist convex bodies M_1 and M_2 such that $V(S \setminus M_1)$ and $V(M_2 \setminus S)$ are arbitrarily close to the first lower bound in (i) and the right-hand side in (ii), respectively. However, this will not be used in the following.

Finally, we provide some rough estimates that will be used repeatedly in the sequel.

Lemma 5.7 *Let Δ_n be a regular simplex inscribed into B^n , and let $\Delta_n^\circ = -n\Delta_n$ be its polar. Then*

- (a) $\ell(\Delta_n) \leq \sqrt{n^3}$, $\ell(\Delta_n^\circ) \leq \sqrt{n}$,
- (b) $V(\Delta_2) \leq 1.3$ and $V(\Delta_n) \leq 1$ for $n \geq 3$,
- (c) $V(\Delta_n^\circ) = n^n V(\Delta_n) \geq (1 + \frac{1}{n})^{\frac{n}{2}} > 1$,
- (d) $V(\Delta_n) \geq n^{-(n+2)} \ell(\Delta_n)$.

Proof. (a) Since $\frac{1}{n}B^n \subset \Delta_n$ and by an application of [66, (7)], we get

$$\ell(\Delta_n) \leq n\ell(B^n) = n \int_{\mathbb{R}^n} \|x\| \gamma_n(dx) = \frac{n^2 \Gamma(\frac{n+1}{2})}{\sqrt{2} \Gamma(\frac{n+2}{2})} \leq \sqrt{n^3}.$$

For (b) and (c), we have

$$\frac{1}{n^n} \leq V(\Delta_n) = \left(1 + \frac{1}{n}\right)^{\frac{n}{2}} \frac{\sqrt{n+1}}{n!} \leq \frac{\sqrt{e} \sqrt{n+1}}{n!} < 1,$$

where the upper bound on the right side only holds for $n \geq 3$.

(d) follows from (a) and (c). \square

6 On the derivatives of the transportation map

Let f and h be probability density functions on \mathbb{R} that are continuous and differentiable on the interiors of their supports, which are assumed to be intervals $I_f, I_h \subset \mathbb{R}$. Then there exists a transportation map $T : I_f \rightarrow I_h$ determined by

$$\int_{-\infty}^x f = \int_{-\infty}^{T(x)} h.$$

For $x \in I_f$ it follows that

$$T'(x) = \frac{f(x)}{h(T(x))}, \tag{15}$$

$$T''(x) = \frac{f(x)^2}{h(T(x))} \left(\frac{f'(x)}{f(x)^2} - \frac{h'(T(x))}{h(T(x))^2} \right). \quad (16)$$

Let g be the standard Gaussian density $g(t) = \sqrt{2\pi}^{-1} e^{-t^2/2}$, $t \in \mathbb{R}$, and for $s \in \mathbb{R}$, let g_s be the truncated Gaussian density

$$g_s(x) = \begin{cases} \left(\int_0^\infty g(t-s) dt \right)^{-1} g(x-s), & \text{if } x \geq 0, \\ 0, & \text{if } x < 0. \end{cases}$$

We frequently use that if $s \geq 0$, then

$$\frac{1}{2} \leq \int_0^\infty g(t-s) dt < 1.$$

We are going to apply (16) either in the case when $h = g$ and $f = g_s$, for some $s \in \mathbb{R}$, or when the roles of f, g are reversed. In particular, we consider the transport maps $\varphi_s : (0, \infty) \rightarrow \mathbb{R}$ and $\psi_s : \mathbb{R} \rightarrow (0, \infty)$ such that

$$\int_0^x g_s = \int_{-\infty}^{\varphi_s(x)} g \quad \text{and} \quad \int_{-\infty}^y g = \int_0^{\psi_s(y)} g_s.$$

Clearly, φ_s and ψ_s are inverses of each other for any given $s \in \mathbb{R}$.

Lemma 6.1 *Let $s \in [0, 0.15]$.*

(i) *If $x \in [0.74, 0.77]$, then $0 < \varphi_s(x) < 0.16$, $1.3 \leq \varphi'_s(x) \leq 2.05$ and $\varphi''_s(x) \leq -0.25$.*

(ii) *If $y \in [0, 0.15]$, then $0 < \psi_s(y) < 0.85$, $0.49 \leq \psi'_s(y) \leq 0.77$ and $\psi''_s(y) \geq 0.07$.*

Proof. We define $\alpha, \beta, \gamma, \delta, \xi > 0$ by the following integrals. The estimates for the values of $\alpha, \beta, \gamma, \delta, \xi > 0$ can be computed numerically.

$$\int_\delta^\infty g = \frac{7}{32}, \quad \text{thus } 0.77 < \delta < 0.78,$$

$$\int_\xi^\infty g = \frac{63}{256}, \quad \text{thus } 0.68 < \xi < 0.69,$$

$$\int_\alpha^\infty g = \frac{1}{4}, \quad \text{thus } 0.67 < \alpha < 0.68,$$

$$\int_\beta^\infty g = \frac{9}{32}, \quad \text{thus } 0.57 < \beta < 0.58,$$

$$\int_\gamma^\infty g = \frac{7}{16}, \quad \text{thus } 0.15 < \gamma < 0.16,$$

and therefore

$$\begin{aligned}\psi_0(0) &= \alpha, \\ \psi_0(\gamma) &= \delta > 0.77,\end{aligned}\tag{17}$$

$$\psi_\gamma(0) = \gamma + \beta < 0.74,\tag{18}$$

$$\psi_\gamma(\gamma) = \gamma + \xi < 0.85.\tag{19}$$

First, we show that if $y \geq 0$, then the map $s \mapsto \psi_s(y) - s$, $s \geq 0$, is strictly decreasing and $\psi_s(y) - s > 0$.

In fact, by definition we have

$$\int_{-\infty}^y g = \int_0^{\psi_s(y)} g_s = \left(\int_{-s}^{\infty} g \right)^{-1} \int_{-s}^{\psi_s(y)-s} g,$$

and hence

$$\int_{-\infty}^y g \int_{-s}^{\infty} g = \int_{-s}^{\infty} g - \int_{\psi_s(y)-s}^{\infty} g$$

or

$$\int_{-s}^{\infty} g \int_y^{\infty} g = \int_{\psi_s(y)-s}^{\infty} g.\tag{20}$$

The left-hand side of (20) is monotone increasing in s , hence the right-hand side is also increasing, which, in turn, implies that $\psi_s(y) - s$ is monotone decreasing as it is in the lower limit of the integral. Moreover, since the left side of (20) is less than $1/2$, it follows that $\psi_s(y) - s > 0$ for $y \geq 0$.

Now, we show that if $y \in [0, \gamma]$, then

$$\psi_s(y) \text{ is a monotone increasing function of } s \geq 0.\tag{21}$$

For the proof of (21), we show that if $0 \leq s < s'$, then the inequality

$$\int_0^{\psi_s(y)} g_{s'} \leq \int_0^{\psi_{s'}(y)} g_{s'} = \int_{-\infty}^y g\tag{22}$$

holds. The inequality (22) implies (21) because, by the positivity of $g_{s'}$, $\psi_{s'}(y) \geq \psi_s(y)$ must hold for

$$\int_0^{\psi_{s'}(y)} g_{s'} = \int_{-\infty}^y g$$

to be true. We set $x := \psi_s(y)$, $\Delta := s' - s \geq 0$ and define

$$A := \int_0^x g(\sigma - s) d\sigma, \quad B := \int_0^{\infty} g(\sigma - s) d\sigma$$

and

$$a := \int_{-\Delta}^0 g(\sigma - s) d\sigma, \quad b := \int_{x-\Delta}^x g(\sigma - s) d\sigma.$$

Note that

$$\int_0^x g_{s'} = \frac{\int_{-\Delta}^{x-\Delta} g(\tau - s) d\tau}{\int_{-\Delta}^{\infty} g(\tau - s) d\tau} = \frac{a + A - b}{a + B}$$

and the right-hand side of (22) equals A/B . Hence (22) is equivalent to

$$\frac{a + A - b}{a + B} \leq \frac{A}{B} \quad \text{or} \quad \frac{a}{b} \leq \frac{1}{1 - \frac{A}{B}}.$$

Since

$$\frac{A}{B} = \int_{-\infty}^y g \geq \frac{1}{2},$$

it is sufficient to show that $a/b \leq 2$.

By the symmetry of g , translation invariance of Lebesgue measure and inserting again $\Delta = s' - s$ and $x = \psi_s(y)$, we get

$$a = \int_s^{s'} g, \quad b = \int_{s-\psi_s(y)}^{s'-\psi_s(y)} g.$$

Thus it remains to be shown that

$$\int_s^{s'} e^{-t^2/2} dt \leq \int_{s-\psi_s(y)}^{s'-\psi_s(y)} 2e^{-t^2/2} dt \quad (23)$$

for $0 \leq s < s'$ and $y \in [0, \gamma]$. To see this, we distinguish two cases.

If $s' - \psi_s(y) \leq 0$, then $2e^{-t^2/2} \geq 1$ for $t \in [s - \psi_s(y), s' - \psi_s(y)] \subset (-\infty, 0]$, since

$$\psi_s(y) - s \leq \psi_s(\gamma) - s \leq \psi_0(\gamma) - 0 = \psi_0(\gamma) < 0.78$$

and $2 \exp(-0.5 \cdot 0.78^2) \geq 1.4 > 1$. Since $e^{-t^2/2} \leq 1$ for $t \in [s, s']$, the assertion follows in this case.

If $s' - \psi_s(y) > 0$, then by the previous reasoning and since $s - \psi_s(y) < 0$, we have

$$\int_s^{\psi_s(y)} e^{-t^2/2} dt \leq \int_{s-\psi_s(y)}^0 2e^{-t^2/2} dt, \quad (24)$$

and since $t \mapsto e^{-t^2/2}$, $t \geq 0$, is decreasing, we have

$$\int_{\psi_s(y)}^{s'} e^{-t^2/2} dt \leq \int_0^{s'-\psi_s(y)} e^{-t^2/2} dt, \quad (25)$$

so that (24) and (25) again imply (23). **Thus, we have proved (22), and, in turn, (21).**

We continue by proving the statements in (ii). We deduce from (17), (18) and (21) that $\psi_s(0) \leq \psi_s(\gamma) < 0.74$, $\psi_s(\gamma) \geq \psi_0(\gamma) > 0.77$, and hence

$$[0.74, 0.77] \subset \psi_s((0, \gamma)) \quad \text{if } s \in [0, \gamma]. \quad (26)$$

We note that if $y \in [0, \gamma]$, then

$$\frac{g'(y)}{g(y)^2} = -\sqrt{2\pi} y e^{y^2/2} \geq -\sqrt{2\pi} \cdot 0.17. \quad (27)$$

On the other hand, if $0 \leq s \leq \gamma$ and $y \geq 0$, then

$$\frac{1}{2} \leq \int_{-s}^{\infty} g \quad \text{and} \quad \psi_s(y) - s \geq \psi_\gamma(y) - \gamma \geq \psi_\gamma(0) - \gamma = \beta > 0.57,$$

and therefore

$$\begin{aligned} \frac{g'_s(\psi_s(y))}{g_s(\psi_s(y))^2} &= -\sqrt{2\pi} \left(\int_{-s}^{\infty} g \right) (\psi_s(y) - s) e^{\frac{(\psi_s(y)-s)^2}{2}} \\ &\leq -\frac{\sqrt{2\pi}}{2} \beta e^{\frac{\beta^2}{2}} < -\sqrt{2\pi} \cdot 0.33. \end{aligned} \quad (28)$$

Combining (27) and (28), for $s, y \in [0, \gamma]$ we get

$$\frac{g'(y)}{g(y)^2} - \frac{g'_s(\psi_s(y))}{g_s(\psi_s(y))^2} \geq \sqrt{2\pi} \cdot 0.15. \quad (29)$$

If $s, y \in [0, \gamma]$, then

$$g_s(\psi_s(y)) \leq \frac{2}{\sqrt{2\pi}} \quad \text{and} \quad g(y) \geq \frac{e^{-\gamma^2/2}}{\sqrt{2\pi}} > \frac{0.98}{\sqrt{2\pi}}. \quad (30)$$

Hence, for $s, y \in [0, \gamma]$ we deduce from (16), (29) and (30) that

$$\psi_s''(y) \geq \frac{0.98^2}{2} \cdot 0.15 > 0.07. \quad (31)$$

In addition, (19) and (21) imply that if $s, y \in [0, \gamma]$, then

$$\psi_s(y) < 0.85. \quad (32)$$

To estimate the first derivative ψ'_s , we use that (15) yields

$$\psi'_s(y) = \frac{g(y)}{g_s(\psi_s(y))}. \quad (33)$$

If $s, y \in [0, \gamma]$, then (30) and (33) yield

$$\psi'_s(y) \geq \frac{0.98/\sqrt{2\pi}}{2/\sqrt{2\pi}} = 0.49. \quad (34)$$

On the other hand, if $s, y \in [0, \gamma]$, then $0 < \psi_s(y) - s \leq \psi_0(y) - 0 \leq \psi_0(\gamma) = \delta < 0.78$, and hence

$$g_s(\psi_s(y)) = \frac{1}{\sqrt{2\pi}} \frac{e^{-(\psi_s(y)-s)^2/2}}{\int_{-s}^{\infty} g} \geq \frac{1}{\sqrt{2\pi}} \frac{e^{-0.78^2/2}}{\int_{-0.16}^{\infty} g} \geq \frac{1}{\sqrt{2\pi}} \cdot 1.3. \quad (35)$$

Hence we deduce from (33) that

$$\psi'_s(y) \leq \frac{1/\sqrt{2\pi}}{1.3/\sqrt{2\pi}} < 0.77. \quad (36)$$

We conclude (ii) from (31), (32), (34) and (36). **This finishes the proof of Lemma 6.1 (ii).**

Finally, we prove part (i) of Lemma 6.1. Turning to φ_s , (26) yields

$$\varphi_s([0.74, 0.77]) \subset (0, \gamma) \quad \text{if } s \in [0, \gamma]. \quad (37)$$

It follows from (29) and (37) that if $s \in [0, \gamma]$ and $x \in [0.74, 0.77]$, then

$$\frac{g'_s(x)}{g_s(x)^2} - \frac{g'(\varphi_s(x))}{g(\varphi_s(x))^2} \leq -\sqrt{2\pi} \cdot 0.15. \quad (38)$$

Now if $s \in [0, \gamma]$ and $x \in [0.74, 0.77]$, then we have

$$g_s(x) \geq \frac{\frac{1}{\sqrt{2\pi}} e^{-0.77^2/2}}{\int_{-0.16}^{\infty} g} > \frac{1.3}{\sqrt{2\pi}}, \quad g(\varphi_s(x)) < \frac{1}{\sqrt{2\pi}}. \quad (39)$$

Hence, (39), (16) and (38) imply that

$$\varphi''_s(x) \leq -1.3^2 \cdot 0.15 < -0.25. \quad (40)$$

To estimate the first derivative φ'_s , we use that (15) yields

$$\varphi'_s(x) = \frac{g_s(x)}{g(\varphi_s(x))}. \quad (41)$$

If $s \in [0, \gamma]$ and $x \in [0.74, 0.77]$, then we conclude from (37) that

$$g(\varphi_s(x)) \geq \frac{e^{-\gamma^2/2}}{\sqrt{2\pi}} > \frac{0.98}{\sqrt{2\pi}} \quad \text{and} \quad g_s(x) \leq \frac{2}{\sqrt{2\pi}},$$

and hence (41) implies that

$$\varphi'_s(x) \leq \frac{2/\sqrt{2\pi}}{0.98/\sqrt{2\pi}} < 2.05. \quad (42)$$

On the other hand, if $s \in [0, \gamma]$ and $x \in [0.74, 0.77]$, then we deduce from (39) and (41) that

$$\varphi'_s(x) > \frac{1.3/\sqrt{2\pi}}{1/\sqrt{2\pi}} = 1.3. \quad (43)$$

We conclude (i) from (40), (37), (42) and (43). \square

In Proposition 6.2, we use the following notation. We fix an $e \in S^n \subset \mathbb{R}^{n+1}$, and identify $e^\perp \subset \mathbb{R}^{n+1}$ with \mathbb{R}^n . For $k \geq n+1$, let $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ be such that

$$\begin{aligned} \sum_{i=1}^k c_i u_i \otimes u_i &= I_n, \\ \sum_{i=1}^k c_i u_i &= o. \end{aligned} \quad (44)$$

For each u_i , we consider

$$\begin{aligned} \tilde{u}_i &= \frac{\sqrt{n}}{\sqrt{n+1}} u_i + \frac{1}{\sqrt{n+1}} e \in S^n, \\ \tilde{c}_i &= \frac{n+1}{n} c_i, \end{aligned} \quad (45)$$

and hence (44) yields that

$$\sum_{i=1}^k \tilde{c}_i \tilde{u}_i \otimes \tilde{u}_i = I_{n+1}.$$

Proposition 6.2 *With the above notation, let $k \leq 2n^2$, let $s \in [0, 0.15]$ and let $\varepsilon \in (0, n^{-56n})$. If*

$$\int_{\mathbb{R}^{n+1}} \prod_{i=1}^k g_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} dx \geq 1 - \varepsilon, \text{ or} \quad (46)$$

$$\int_{\mathbb{R}^{n+1}}^* \sup_{x = \sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k g_s(\theta_i)^{\tilde{c}_i} dx \leq 1 + \varepsilon, \quad (47)$$

then there exists a regular simplex with vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ and $i_1 < \dots < i_{n+1}$ such that $\angle(u_{i_j}, w_j) < n^{14n} \varepsilon^{1/4}$ for $j = 1, \dots, n+1$.

Proof. According to Lemma 4.4, we may assume

$$\tilde{c}_1 \cdots \tilde{c}_{n+1} \det[\tilde{u}_1, \dots, \tilde{u}_{n+1}]^2 \geq \binom{k}{n+1}^{-1}. \quad (48)$$

For $\eta = n^{10n} \varepsilon^{1/4} < 1$, we claim that if $i \in \{1, \dots, k\}$, then

$$\tilde{c}_i \leq \eta^2, \text{ or there exists some } j \in \{1, \dots, n+1\} \text{ with } \angle(\tilde{u}_i, \tilde{u}_j) \leq \eta. \quad (49)$$

We suppose that (49) does not hold, hence we may assume

$$\tilde{c}_{n+2} > \eta^2 \text{ and } \angle(\tilde{u}_i, \tilde{u}_{n+2}) > \eta \text{ for } i = 1, \dots, n+1.$$

We can write $\tilde{u}_{n+2} = \sum_{i=1}^{n+1} \lambda_i \tilde{u}_i$, where $\lambda_1, \dots, \lambda_{n+1} \in \mathbb{R}$ are uniquely determined and satisfy $\lambda_1 + \dots + \lambda_{n+1} = 1$. Hence we may assume that $\lambda_1 \geq \frac{1}{n+1}$. Therefore $\tilde{c}_{n+2} > \eta^2$, $\tilde{c}_1 \leq 1$ and (48) imply

$$\tilde{c}_2 \cdots \tilde{c}_{n+2} \det[\tilde{u}_2, \dots, \tilde{u}_{n+2}]^2 \geq \binom{k}{n+1}^{-1} \frac{\eta^2}{(n+1)^2} \geq \frac{(n+1)!}{(2n^2)^{n+1}} \frac{\eta^2}{(n+1)^2}.$$

Here $\frac{(n+1)!}{(n+1)^2} \geq \frac{n^n}{(n+1)e^n} > \frac{n^{n-1}}{3^{n+1}}$, and thus

$$\tilde{c}_2 \cdots \tilde{c}_{n+2} \det[\tilde{u}_2, \dots, \tilde{u}_{n+2}]^2 \geq \frac{\eta^2}{3^{n+1}2^{n+1}\eta^{n+3}} > \frac{\eta^2}{n^{4n+6}}. \quad (50)$$

In addition, $\angle(\tilde{u}_1, \tilde{u}_{n+2}) > \eta$ yields

$$\|\tilde{u}_1 - \tilde{u}_{n+2}\| > \eta/2. \quad (51)$$

We prove (49) separately for (46) and (47).

We start with the Brascamp-Lieb inequality; namely, we assume that (46) holds. We observe that if $i = 1, \dots, k$, then

$$0.74 < \langle x, \tilde{u}_i \rangle < 0.77 \quad \text{for } x \in 0.755\sqrt{n+1}e + 0.01B^n,$$

which can be checked by directly computing the inner product $\langle x, \tilde{u}_i \rangle$ using the definition of the vectors \tilde{u}_i in (45).

Define

$$\Xi := 0.755\sqrt{n+1}e + 0.005 \frac{\tilde{u}_1 - \tilde{u}_{n+2}}{\|\tilde{u}_1 - \tilde{u}_{n+2}\|} + 0.001B^n \subset 0.755\sqrt{n+1}e + 0.01B^n.$$

It follows using also (51), $\langle e, \tilde{u}_1 - \tilde{u}_{n+2} \rangle = 0$ and $V(B^n) = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2}+1)} > \frac{(2e\pi)^{\frac{n}{2}}}{n^{n/2}\sqrt{2\pi}e^{1/(6n)}} > \frac{1}{n^{n/2}}$ that

$$\langle x, \tilde{u}_1 \rangle, \dots, \langle x, \tilde{u}_k \rangle \in [0.74, 0.77] \quad \text{for } x \in \Xi, \quad (52)$$

$$\langle x, \tilde{u}_1 \rangle - \langle x, \tilde{u}_{n+2} \rangle \geq 0.002\eta > 2^{-9}\eta \quad \text{for } x \in \Xi, \quad (53)$$

$$\Xi \subset \mathcal{C} := \{x \in \mathbb{R}^{n+1} : \langle \tilde{u}_i, x \rangle > 0 \forall i = 1, \dots, k\}, \quad (54)$$

$$V(\Xi) = 0.001^n V(B^n) > \frac{1}{n^{11n}}, \quad (55)$$

where (54) is a consequence of (52). In addition, we consider the map $\Theta : \mathcal{C} \rightarrow \mathbb{R}^{n+1}$ with

$$\Theta(x) = \sum_{i=1}^k \tilde{c}_i \varphi_s(\langle \tilde{u}_i, x \rangle) \tilde{u}_i, \quad x \in \mathcal{C},$$

which satisfies

$$d\Theta(x) = \sum_{i=1}^k \tilde{c}_i \varphi'_s(\langle \tilde{u}_i, x \rangle) \tilde{u}_i \otimes \tilde{u}_i.$$

As we have seen, $d\Theta$ is positive definite and $\Theta : \mathcal{C} \rightarrow \mathbb{R}^{n+1}$ is injective (see [9, 10]). Therefore, applying first (46), then (12), and after that the definition of Θ and (10), we obtain

$$1 - \varepsilon \leq \int_{\mathbb{R}^{n+1}} \prod_{i=1}^k g_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} dx = \int_{\mathcal{C}} \prod_{i=1}^k g_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} dx$$

$$\begin{aligned}
&\leq \int_{\mathcal{C}} \left(\prod_{i=1}^k g(\varphi_s(\langle x, \tilde{u}_i \rangle))^{\tilde{c}_i} \right) \left(\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \right) dx \\
&= \left(\frac{1}{2\pi} \right)^{\frac{n+1}{2}} \int_{\mathcal{C}} \left(\prod_{i=1}^k e^{-\tilde{c}_i \varphi_s(\langle x, \tilde{u}_i \rangle)^2/2} \right) \left(\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \right) dx \\
&\leq \left(\frac{1}{2\pi} \right)^{\frac{n+1}{2}} \int_{\mathcal{C}} e^{-\|\Theta(x)\|^2/2} \left(\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \right) dx. \tag{56}
\end{aligned}$$

We deduce from (9) that

$$\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \leq \det \left(\sum_{i=1}^k \tilde{c}_i \varphi'_s(\langle x, \tilde{u}_i \rangle) \tilde{u}_i \otimes \tilde{u}_i \right) = \det(d\Theta(x)) \tag{57}$$

for any $x \in \mathcal{C}$.

If $s \in [0, 0.15]$ and $x \in \Xi$, then we can improve (57) using Corollary 4.3, based on (48) and (50) with

$$\beta_0 = \frac{\eta^2}{n^{4n+6}}.$$

Hence, applying first Corollary 4.3, then Lemma 6.1 (i), (52), (53) and finally $\eta < 1$, we get

$$\begin{aligned}
\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} &\leq \left(1 + \frac{\beta_0 (\varphi'_s(\langle x, \tilde{u}_1 \rangle) - \varphi'_s(\langle x, \tilde{u}_{n+2} \rangle))^2}{4(\varphi'_s(\langle x, \tilde{u}_1 \rangle) + \varphi'_s(\langle x, \tilde{u}_{n+2} \rangle))^2} \right)^{-1} \det(d\Theta(x)) \\
&\leq \left(1 + \frac{\beta_0 (0.25(\langle x, \tilde{u}_1 \rangle - \langle x, \tilde{u}_{n+2} \rangle))^2}{4(2 \cdot 2.05)^2} \right)^{-1} \det(d\Theta(x)) \\
&\leq \left(1 + \frac{\eta^4 0.25^2 2^{-18}}{n^{4n+6} 16 \cdot 2.05^2} \right)^{-1} \det(d\Theta(x)) \\
&\leq \left(1 + \frac{\eta^4}{n^{4n+35}} \right)^{-1} \det(d\Theta(x)) \leq \left(1 - \frac{\eta^4}{n^{4n+36}} \right) \det(d\Theta(x)).
\end{aligned}$$

Moreover, if $s \in [0, 0.15]$ and $x \in \Xi$ we deduce from (57) and Lemma 6.1 (i) that

$$\det(d\Theta(x)) \geq \prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \geq \prod_{i=1}^k 1^{\tilde{c}_i} = 1.$$

Thus if $s \in [0, 0.15]$ and $x \in \Xi$, then

$$\prod_{i=1}^k \varphi'_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} \leq \det(d\Theta(x)) - \frac{\eta^4}{n^{4n+36}}. \tag{58}$$

In addition, if $s \in [0, 0.15]$ and $x \in \Xi$, then $\langle x, \tilde{u}_i \rangle \in [0.74, 0.77]$ by (52) and hence $\varphi_s(\langle x, \tilde{u}_i \rangle) \subset (0, \gamma)$ by (37). Therefore, the definition of $\Theta(x)$, (10) and (11) imply

$$\|\Theta(x)\|^2 \leq \sum_{i=1}^k \tilde{c}_i \varphi_s(\langle \tilde{u}_i, x \rangle)^2 \leq \sum_{i=1}^k \tilde{c}_i 0.16^2 = 0.16^2(n+1),$$

and hence

$$\left(\frac{1}{2\pi}\right)^{\frac{n+1}{2}} e^{-\|\Theta(x)\|^2/2} \geq \left(\frac{1}{2\pi}\right)^{\frac{n+1}{2}} e^{-(n+1)0.16^2/2} > n^{-n-3}.$$

Applying first (54), using (57) and (58) in (56), then the substitution $z = \Theta(x)$, and finally also (55), we get

$$\begin{aligned} 1 - \varepsilon &\leq \left(\frac{1}{2\pi}\right)^{\frac{n+1}{2}} \int_{\mathcal{C}} e^{-\|\Theta(x)\|^2/2} \det(d\Theta(x)) dx - \int_{\Xi} \frac{\eta^4}{n^{5n+39}} dx \\ &\leq \left(\frac{1}{2\pi}\right)^{\frac{n+1}{2}} \int_{\mathbb{R}^{n+1}} e^{-\|z\|^2/2} dz - \frac{1}{n^{11n}} \frac{\eta^4}{n^{5n+39}} \leq 1 - \frac{\eta^4}{n^{39n}}. \end{aligned}$$

This contradicts $\eta = n^{10n} \varepsilon^{1/4}$, and hence we conclude (49) in the case of the Brascamp-Lieb inequality.

Now we consider the reverse Brascamp-Lieb inequality; namely, we assume that (47) holds. We observe that if $i \in \{1, \dots, k\}$, then

$$0 \leq \langle x, \tilde{u}_i \rangle \leq 0.15 \quad \text{for } x \in 0.1\sqrt{n+1}e + 0.05B^n,$$

and define

$$\tilde{\Xi} := 0.1\sqrt{n+1}e + 0.03 \frac{\tilde{u}_1 - \tilde{u}_{n+2}}{\|\tilde{u}_1 - \tilde{u}_{n+2}\|} + 0.01B^n \subset 0.1\sqrt{n+1}e + 0.05B^n.$$

It follows using again (51), $\langle e, \tilde{u}_1 - \tilde{u}_{n+2} \rangle = 0$ and $V(B^n) > \frac{1}{n^{n/2}}$ that

$$\langle y, \tilde{u}_1 \rangle, \dots, \langle y, \tilde{u}_k \rangle \in [0, 0.15] \quad \text{for } y \in \tilde{\Xi} \tag{59}$$

$$\langle y, \tilde{u}_1 \rangle - \langle y, \tilde{u}_{n+2} \rangle \geq 0.01\eta > 2^{-7}\eta \quad \text{for } y \in \tilde{\Xi} \tag{60}$$

$$V(\tilde{\Xi}) = 0.01^n V(B^n) > \frac{1}{n^{8n}}. \tag{61}$$

In addition, we consider the map $\Psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ with

$$\Psi(y) = \sum_{i=1}^k \tilde{c}_i \psi_s(\langle \tilde{u}_i, y \rangle) \tilde{u}_i,$$

which satisfies

$$d\Psi(y) = \sum_{i=1}^k \tilde{c}_i \psi'_s(\langle \tilde{u}_i, y \rangle) \tilde{u}_i \otimes \tilde{u}_i.$$

As we have seen, $d\Psi$ is positive definite and $\Psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ is injective (see [9, 10]). Therefore, applying first (47), then the definition of Ψ imply

$$\begin{aligned} 1 + \varepsilon &\geq \int_{\mathbb{R}^{n+1}}^* \sup_{x=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k g_s(\theta_i)^{\tilde{c}_i} dx \\ &\geq \int_{\mathbb{R}^{n+1}}^* \left(\sup_{\Psi(y)=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k g_s(\theta_i)^{\tilde{c}_i} \right) \det(d\Psi(y)) dy \\ &\geq \int_{\mathbb{R}^{n+1}} \left(\prod_{i=1}^k g_s(\psi_s(\langle \tilde{u}_i, y \rangle))^{\tilde{c}_i} \right) \det \left(\sum_{i=1}^k \tilde{c}_i \psi'_s(\langle \tilde{u}_i, y \rangle) \tilde{u}_i \otimes \tilde{u}_i \right) dy. \end{aligned} \quad (62)$$

Using (9), for $y \in \mathbb{R}^{n+1}$ we can bound the determinant in (62) from below by

$$\det \left(\sum_{i=1}^k \tilde{c}_i \psi'_s(\langle y, \tilde{u}_i \rangle) \tilde{u}_i \otimes \tilde{u}_i \right) \geq \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i}. \quad (63)$$

If $s \in [0, 0.15]$ and $y \in \tilde{\Xi}$, an application of Corollary 4.3 with $\beta_0 = \eta^2/n^{4n+6}$, Lemma 6.1 (ii), (59) and (60), allow us to improve (63) to get

$$\begin{aligned} \det \left(\sum_{i=1}^k \tilde{c}_i \psi'_s(\langle y, \tilde{u}_i \rangle) \tilde{u}_i \otimes \tilde{u}_i \right) &\geq \left(1 + \frac{\beta_0 (\psi'_s(\langle y, \tilde{u}_1 \rangle) - \psi'_s(\langle y, \tilde{u}_{n+2} \rangle))^2}{4(\psi'_s(\langle y, \tilde{u}_1 \rangle) + \psi'_s(\langle y, \tilde{u}_{n+2} \rangle))^2} \right) \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i} \\ &\geq \left(1 + \frac{\beta_0 (0.07(\langle y, \tilde{u}_1 \rangle - \langle y, \tilde{u}_{n+2} \rangle))^2}{4(2 \cdot 0.77)^2} \right) \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i} \\ &\geq \left(1 + \frac{\eta^4 0.07^2 2^{-14}}{n^{4n+6} 16 \cdot 0.77^2} \right) \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i} \\ &\geq \left(1 + \frac{\eta^4}{n^{4n+31}} \right) \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i}. \end{aligned}$$

Moreover, if $s \in [0, 0.15]$ and $y \in \tilde{\Xi}$ we deduce from Lemma 6.1 (ii) and (11) that

$$\prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i} \geq \prod_{i=1}^k 0.49^{\tilde{c}_i} = 0.49^{n+1} > n^{-2n}.$$

Thus if $s \in [0, 0.15]$ and $y \in \tilde{\Xi}$, then

$$\det \left(\sum_{i=1}^k \tilde{c}_i \psi'_s(\langle y, \tilde{u}_i \rangle) \tilde{u}_i \otimes \tilde{u}_i \right) \geq \prod_{i=1}^k \psi'_s(\langle y, \tilde{u}_i \rangle)^{\tilde{c}_i} + \frac{\eta^4}{n^{6n+31}}. \quad (64)$$

Further in (62), if $s \in [0, 0.15]$ and $y \in \tilde{\Xi}$, then (59), Lemma 6.1 (ii), (11) and (35) imply

$$\prod_{i=1}^k g_s(\psi_s(\langle \tilde{u}_i, y \rangle))^{\tilde{c}_i} \geq \left(\frac{1.3}{\sqrt{2\pi}} \right)^{n+1} \geq 2^{-n-1} \geq n^{-n-1}. \quad (65)$$

Applying first (63), (64) and (65) in (62), and then (12) and (61), we deduce that if $s \in [0, 0.15]$, then

$$\begin{aligned} 1 + \varepsilon &\geq \int_{\mathbb{R}^{n+1}} \left(\prod_{i=1}^k g_s(\psi_s(\langle \tilde{u}_i, y \rangle))^{\tilde{c}_i} \right) \left(\prod_{i=1}^k \psi'_s(\langle \tilde{u}_i, y \rangle)^{\tilde{c}_i} \right) dy + \int_{\tilde{\Xi}} \frac{\eta^4}{n^{7n+32}} dy \\ &\geq \int_{\mathbb{R}^{n+1}} \left(\prod_{i=1}^k g(\langle \tilde{u}_i, y \rangle)^{\tilde{c}_i} \right) dy + \frac{\eta^4}{n^{15n+31}} \\ &\geq \left(\frac{1}{\sqrt{2\pi}} \right)^{\frac{n+1}{2}} \int_{\mathbb{R}^{n+1}} e^{-\|y\|^2/2} dy + \frac{\eta^4}{n^{31n}} = 1 + \frac{\eta^4}{n^{31n}}. \end{aligned}$$

This contradicts $\eta = n^{10n}\varepsilon^{1/4}$, and hence we conclude (49) also in the case of the reverse Brascamp-Lieb inequality.

Now we return to the proof of Proposition 6.2. Since $k \leq 2n^2$ and $\varepsilon < n^{-56n}$, we have $\eta < 1/(6kn)$. Since (49) is available now, we can apply Corollary 5.3, which yields the existence of an orthonormal basis $\tilde{w}_1, \dots, \tilde{w}_{n+1}$ of \mathbb{R}^{n+1} such that $\langle e, \tilde{w}_i \rangle = \frac{1}{\sqrt{n+1}}$ and $\|\tilde{u}_i - \tilde{w}_i\| \leq \angle(\tilde{u}_i, \tilde{w}_i) < 6n^3\eta$ for $\eta = n^{10n}\varepsilon^{1/4}$ and $i = 1, \dots, n+1$. Now we consider the vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ of the regular simplex which are defined by the relations $\tilde{w}_i = \sqrt{\frac{n}{n+1}} w_i + \sqrt{\frac{1}{n+1}} e$ for $i = 1, \dots, n+1$. Therefore

$$\angle(u_i, w_i) < \frac{\pi}{2} \|u_i - w_i\| = \frac{\pi}{2} \sqrt{\frac{n+1}{n}} \|\tilde{u}_i - \tilde{w}_i\| < 18n^3\eta < n^{14n}\varepsilon^{1/4}$$

for $i = 1, \dots, n+1$. In turn, we conclude Proposition 6.2. \square

We will actually use the Brascamp-Lieb inequality and its reverse for the function

$$\tilde{g}_s(t) = \mathbf{1}\{t \geq 0\} \exp\left(-\frac{(t-s)^2}{2}\right) \quad (66)$$

for $s \in \mathbb{R}$, i.e.

$$\tilde{g}_s = \left(\int_{\mathbb{R}} \tilde{g}_s \right) g_s.$$

We note that if $s \geq 0$, then

$$\int_{\mathbb{R}} \tilde{g}_s \geq \frac{\sqrt{2\pi}}{2} > 1. \quad (67)$$

From Proposition 6.2 and (11) we deduce the following strengthened version of the Brascamp-Lieb inequality and its reverse for \tilde{g}_s .

Corollary 6.3 *Using the same notation as in Proposition 6.2, let $k \leq 2n^2$, let $s \in [0, 0.15]$ and let $\varrho \in (0, 1)$.*

If for any regular simplex with vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ and any subset $\{i_1, \dots, i_{n+1}\} \subset \{1, \dots, k\}$, there exists $j \in \{1, \dots, n+1\}$ such that $\angle(u_{i_j}, w_j) \geq \varrho$, then

$$\int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle x, \tilde{u}_i \rangle)^{\tilde{c}_i} dx \leq (1 - n^{-56n} \varrho^4) \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1},$$

$$\int_{\mathbb{R}^{n+1}} \sup_{x = \sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k \tilde{g}_s(\theta_i)^{\tilde{c}_i} dx \geq (1 + n^{-56n} \varrho^4) \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1}.$$

7 An almost regular simplex for Theorem 1.3 and Theorem 1.6 (a), (b)

The entire section is devoted to proving the following statement.

Proposition 7.1 *Let $n+1 \leq k \leq 2n^2$, $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ be such that*

$$\sum_{i=1}^k c_i u_i \otimes u_i = I_n,$$

$$\sum_{i=1}^k c_i u_i = o,$$

and $\ell(C) \geq (1 - \varepsilon)\ell(\Delta_n)$ holds for $C = \text{conv}\{u_1, \dots, u_k\}$ and $\varepsilon \in (0, n^{-60n})$.

Then for $\eta = n^{15n} \varepsilon^{\frac{1}{4}} \in (0, 1)$, there exists a regular simplex with vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ and $\{i_1, \dots, i_{n+1}\} \subset \{1, \dots, k\}$ such that

$$\angle(u_{i_j}, w_j) \leq \eta \quad \text{for } j = 1, \dots, n+1.$$

We fix $e \in S^n \subset \mathbb{R}^{n+1}$, and identify $e^\perp \subset \mathbb{R}^{n+1}$ with \mathbb{R}^n . As before Proposition 6.2, for each u_i , we consider

$$\tilde{u}_i = \frac{\sqrt{n}}{\sqrt{n+1}} u_i + \frac{1}{\sqrt{n+1}} e \in S^n,$$

$$\tilde{c}_i = \frac{n+1}{n} c_i,$$

and hence

$$\sum_{i=1}^k \tilde{c}_i \tilde{u}_i \otimes \tilde{u}_i = I_{n+1}, \quad (68)$$

$$\sum_{i=1}^k \tilde{c}_i = n + 1. \quad (69)$$

Indirectly, we assume that Proposition 7.1 does not hold, and we aim at a contradiction. We deduce from Corollary 6.3 and (67) that if $s \in [0, 0.15]$, then

$$\int_{\mathbb{R}^{n+1}}^* \sup_{y=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k \tilde{g}_s(\theta_i)^{\tilde{c}_i} dy \geq (1 + n^{-56n} \eta^4) \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} > \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} + n^{-56n} \eta^4. \quad (70)$$

Next we provide the following general auxiliary result (which holds independently of the indirect assumption).

Lemma 7.2 *If $s \in \mathbb{R}$ and C is defined as above, then*

$$\begin{aligned} \text{(i)} \quad & (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n(r\sqrt{n}C) dr \geq \int_{\mathbb{R}^{n+1}}^* \sup_{y=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k \tilde{g}_s(\theta_i)^{\tilde{c}_i} dy, \\ \text{(ii)} \quad & (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n(r\sqrt{n}\Delta_n) dr = \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1}. \end{aligned}$$

Proof. We consider the convex cone

$$\begin{aligned} \mathcal{C}_0 &:= \left\{ \sum_{i=1}^k \xi_i \tilde{u}_i : \xi_i \geq 0 \text{ for } i = 1, \dots, k \right\} \\ &= \left\{ \sum_{i=1}^k r\sqrt{n}\lambda_i u_i + re : r \geq 0, \lambda_i \in [0, 1], \sum_{i=1}^k \lambda_i = 1 \right\}. \end{aligned}$$

Then we clearly have $x + re \in \mathcal{C}_0$ for $x \in \mathbb{R}^n$ and $r \in \mathbb{R}$ if and only if $r \geq 0$ and $x \in r\sqrt{n}C$. If $y = \sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i$ for $\theta_1, \dots, \theta_k \geq 0$, then $\langle y, e \rangle = (\sum_{i=1}^k \tilde{c}_i \theta_i) / \sqrt{n+1}$, and hence we deduce from (10), (68) and (69) that

$$\begin{aligned} \int_{\mathbb{R}^{n+1}}^* \sup_{y=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i} \prod_{i=1}^k \tilde{g}_s(\theta_i)^{\tilde{c}_i} dy &= \int_{\mathcal{C}_0}^* \sup_{y=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{u}_i, \theta_i \geq 0} e^{-\frac{1}{2} \sum_{i=1}^k \tilde{c}_i \theta_i^2 + s \sum_{i=1}^k \tilde{c}_i \theta_i - \frac{s^2}{2} \sum_{i=1}^k \tilde{c}_i} dy \\ &\leq e^{-\frac{(n+1)s^2}{2}} \int_{\mathcal{C}_0} e^{-\frac{1}{2} \|y\|^2 + s \langle y, e \rangle \sqrt{n+1}} dy \\ &= e^{-\frac{(n+1)s^2}{2}} \int_0^\infty \int_{r\sqrt{n}C} e^{-\frac{1}{2} (\|x\|^2 + r^2) + sr\sqrt{n+1}} dx dr \\ &= (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n(r\sqrt{n}C) dr, \end{aligned}$$

thus we have obtained (i).

For (ii), let w_1, \dots, w_{n+1} be the vertices of Δ_n , and let

$$\tilde{w}_i = \frac{\sqrt{n}}{\sqrt{n+1}} w_i + \frac{1}{\sqrt{n+1}} e$$

for $i = 1, \dots, n+1$. Then $\tilde{w}_1, \dots, \tilde{w}_{n+1}$ form an orthonormal basis of \mathbb{R}^{n+1} , and hence

$$\sum_{i=1}^{n+1} \tilde{w}_i \otimes \tilde{w}_i = I_{n+1}$$

(with $\tilde{c}_i = 1$ in this case). Moreover for any $y \in \mathbb{R}^{n+1}$, there exist unique $\theta_1, \dots, \theta_{n+1} \in \mathbb{R}$ satisfying $y = \sum_{i=1}^{n+1} \theta_i \tilde{w}_i$, in fact, we have $\theta_i = \langle y, \tilde{w}_i \rangle$ and $\sum_{i=1}^{n+1} \theta_i^2 = \|y\|^2$ for $i = 1, \dots, n+1$. By the preceding argument, we deduce

$$\begin{aligned} (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n(r\sqrt{n}\Delta_n) dr &= \int_{\mathbb{R}^{n+1}}^* \sup_{y=\sum_{i=1}^k \theta_i \tilde{w}_i} \prod_{i=1}^{n+1} \tilde{g}_s(\theta_i) dy \\ &= \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1}, \end{aligned}$$

where we used Fubini's theorem for the second equality. \square

We apply the change of parameter $\tau = s\sqrt{n(n+1)}$ and substitution $t = r\sqrt{n}$ in Lemma 7.2, and conclude with the help of the reverse Brascamp-Lieb inequality (14) that if $\tau \in \mathbb{R}$, then

$$\begin{aligned} \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} \gamma_n(tC) dt &= e^{-\frac{s^2(n+1)}{2}} \sqrt{n} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n(r\sqrt{n}C) dr \\ &\geq \frac{\sqrt{n}}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^{n+1}}^* \sup_{y=\sum_{i=1}^k \tilde{c}_i \theta_i \tilde{w}_i} \prod_{i=1}^k \tilde{g}_s(\theta_i) \tilde{c}_i dy \\ &\geq \frac{\sqrt{n}}{(2\pi)^{\frac{n}{2}}} \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} \\ &= \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} \gamma_n(t\Delta_n) dt. \end{aligned}$$

Hence, we get

$$\int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(t\Delta_n)) dt \geq \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(tC)) dt. \quad (71)$$

In addition, if $\tau \in [0, 0.15n] \subset [0, 0.15\sqrt{n(n+1)}]$, so that $s = \tau/\sqrt{n(n+1)} \in [0, 0.15)$, then using (70), instead of the reverse Brascamp-Lieb inequality (14), we obtain

$$\int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} \gamma_n(tC) dt \geq \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} \gamma_n(t\Delta_n) dt + \frac{\sqrt{n}}{(2\pi)^{\frac{n}{2}}} n^{-56n} \eta^4,$$

and therefore

$$\int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(t\Delta_n)) dt \geq \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(tC)) dt + \frac{\sqrt{n}}{(2\pi)^{\frac{n}{2}}} n^{-56n} \eta^4. \quad (72)$$

Integrating (71) for $\tau \in \mathbb{R} \setminus [0, 0.15n]$ and (72) for $\tau \in [0, 0.15n]$, we deduce that

$$\begin{aligned} & \int_{-\infty}^\infty \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(t\Delta_n)) dt d\tau \\ & \geq \int_{-\infty}^\infty \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(tC)) dt d\tau + 0.15n \frac{\sqrt{n}}{(2\pi)^{\frac{n}{2}}} n^{-56n} \eta^4 \end{aligned} \quad (73)$$

Since for any $t \in \mathbb{R}$, we have

$$\int_{-\infty}^\infty e^{-\frac{1}{2n}(t-\tau)^2} d\tau = \sqrt{2\pi n}$$

we deduce from (2) and (73) that

$$\begin{aligned} \ell(C) &= \int_0^\infty (1 - \gamma_n(tC)) dt \\ &= \frac{1}{\sqrt{2\pi n}} \int_{-\infty}^\infty \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(tC)) dt d\tau \\ &\leq \frac{1}{\sqrt{2\pi n}} \int_{-\infty}^\infty \int_0^\infty e^{-\frac{1}{2n}(t-\tau)^2} (1 - \gamma_n(t\Delta_n)) dt d\tau - \frac{0.15n}{(2\pi)^{\frac{n+1}{2}}} n^{-56n} \eta^4 \\ &= \ell(\Delta_n) - \frac{0.15n}{(2\pi)^{\frac{n+1}{2}}} n^{-56n} \eta^4. \end{aligned} \quad (74)$$

Hence, Lemma 5.7 (a), (74) and the hypothesis yield

$$(1 - \varepsilon)\ell(\Delta_n) \leq \ell(C) < (1 - n^{-60n}\eta^4)\ell(\Delta_n).$$

This contradicts $\eta = n^{15n}\varepsilon^{\frac{1}{4}}$, and in turn implies Proposition 7.1.

8 Proof of Theorem 1.3 and of Theorem 1.6 (a), (b)

For Theorem 1.6, let μ be a centered isotropic measure on S^{n-1} , and let $K := Z_\infty(\mu)$, and hence $\text{supp } \mu = \partial K \cap S^{n-1}$. In particular, under the assumptions of Theorem 1.3 and of Theorem 1.6 (a), we have $\ell(K) \geq (1 - \varepsilon)\ell(\Delta_n)$. First, we assume that

$$0 < \varepsilon < n^{-100n}.$$

It follows from Lemma 2.1 and John's theorem that there exist $k \geq n + 1$ with $k \leq 2n^2$, $u_1, \dots, u_k \in \partial K \cap S^{n-1}$ and $c_1, \dots, c_k > 0$ such that

$$\begin{aligned} \sum_{i=1}^k c_i u_i \otimes u_i &= I_n, \\ \sum_{i=1}^k c_i u_i &= o. \end{aligned}$$

We write μ_0 to denote the centered discrete isotropic measure with $\text{supp } \mu_0 = \{u_1, \dots, u_k\}$ and $\mu_0(\{u_i\}) = c_i$ for $i = 1, \dots, k$, and define

$$C := Z_\infty(\mu_0) = \text{conv}\{u_1, \dots, u_k\}.$$

Since $\ell(C) \geq \ell(K) \geq (1 - \varepsilon)\ell(\Delta_n)$ and $0 < \varepsilon < n^{-60n}$, it follows from Proposition 7.1 that we may assume that the vertices w_1, \dots, w_{n+1} of Δ_n satisfy

$$\angle(u_i, w_i) \leq \eta \text{ for } \eta = n^{15n}\varepsilon^{\frac{1}{4}} \quad \text{and} \quad i = 1, \dots, n+1. \quad (75)$$

For the simplex

$$S_0 = \text{conv}\{u_1, \dots, u_{n+1}\} \subset K,$$

we deduce from (75) and Lemma 5.4 (where we use $\eta < 1/(2n)$) that $S_0^\circ \subset (1 + 2n\eta)\Delta_n^\circ$, and hence

$$\tilde{\Delta}_n := (1 + 2n\eta)^{-1}\Delta_n \subset S_0 \subset K. \quad (76)$$

We note that

$$\begin{aligned} 0 < \ell(\tilde{\Delta}_n) - \ell(\Delta_n) &= \int_{\mathbb{R}^n} \|x\|_{\tilde{\Delta}_n} d\gamma_n(x) - \ell(\Delta_n) \\ &\leq (1 + 2n\eta)\ell(\Delta_n) - \ell(\Delta_n) = 2n\eta \ell(\Delta_n). \end{aligned} \quad (77)$$

Proof of Theorem 1.3: Let $\xi > 0$ be minimal such that

$$K \subset (1 + \xi)\tilde{\Delta}_n = (1 + \xi)(1 + 2n\eta)^{-1}\Delta_n.$$

Then Lemma 5.6 and Lemma 5.7 (d) imply that

$$V(K \setminus \tilde{\Delta}_n) \geq \frac{\xi}{n+1} V(\tilde{\Delta}_n) = \frac{\xi}{(n+1)(1+2n\eta)^n} V(\Delta_n) > \frac{\xi}{n^{2n+4}} \ell(\Delta_n). \quad (78)$$

It follows from $K \subset B^n$, (76) and (78) that

$$\begin{aligned} \gamma_n(tK) &\geq \gamma_n(t\tilde{\Delta}_n) \text{ for } t > 0, \text{ and} \\ \gamma_n(tK) &\geq \gamma_n(t\tilde{\Delta}_n) + \frac{e^{-\frac{1}{2}}}{(2\pi)^{\frac{n}{2}}} \frac{t^n \xi}{n^{2n+4}} \ell(\Delta_n) \text{ for } t \in (0, 1], \end{aligned}$$

where we used that $e^{-\frac{t^2}{2}} \geq e^{-\frac{1}{2}}$ for $t \in (0, 1]$, and in turn we deduce from (2) that

$$\ell(\tilde{\Delta}_n) - \ell(K) \geq \int_0^1 \gamma_n(tK) - \gamma_n(t\tilde{\Delta}_n) dt \geq \int_0^1 \frac{t^n \xi}{n^{4n+4}} t \ell(\Delta_n) dt > n^{-(4n+5)} \ell(\Delta_n) \xi.$$

We conclude from (77) that

$$(1 - \varepsilon)\ell(\Delta_n) \leq \ell(K) \leq (1 - n^{-(4n+5)}\xi + 2n\eta)\ell(\Delta_n),$$

and hence $\eta = n^{15n} \varepsilon^{\frac{1}{4}}$ implies

$$\xi \leq n^{4n+5}(2n\eta + \varepsilon) < n^{23n} \varepsilon^{\frac{1}{4}}.$$

It follows from (76) and the definition of ξ that

$$(1 - 2n\eta)\Delta_n \subset K \subset (1 + \xi)\Delta_n. \quad (79)$$

Since $\Delta_n \subset B^n$, $\eta = n^{15n} \varepsilon^{\frac{1}{4}} < n^{23n} \varepsilon^{\frac{1}{4}} =: \tilde{\xi}$ and $\xi < \tilde{\xi}$, we conclude for the Hausdorff distance that $\delta_H(K, \Delta_n) < n^{23n} \varepsilon^{\frac{1}{4}}$.

To estimate the symmetric difference distance of K and Δ_n , Lemma 5.7 (b), (79) and $\xi < \tilde{\xi} \leq n^{-2n}$ yield

$$\delta_{\text{vol}}(K, \Delta_n) \leq \left((1 + \tilde{\xi})^n - (1 - \tilde{\xi})^n \right) V(\Delta_n) \leq 2\tilde{\xi}n(1 + \tilde{\xi})^{n-1}V(\Delta_n) < n^{25n} \varepsilon^{\frac{1}{4}},$$

which finishes the proof of Theorem 1.3 if $\varepsilon < n^{-100n}$. However, if $\varepsilon \geq n^{-100n}$, then Theorem 1.3 trivially holds as $\delta_{\text{vol}}(M, \Delta_n) < \kappa_n$ and $\delta_H(M, \Delta_n) < 1$ for any convex body $M \subset B^n$ by the choice of the constant $c = n^{26n}$. \square

Proof of Theorem 1.6 (a), (b): We assume that $\ell(Z_\infty(\mu)) \geq (1 - \varepsilon)\ell(\Delta_n)$ is available.

Let $\alpha_0 = 9 \cdot 2^{n+2}n^{2n+2}$ be the constant of Lemma 5.5. If for any $u \in \text{supp } \mu$ there exists a w_i such that $\angle(u, w_i) \leq \alpha_0\eta$, then

$$\delta_H(\text{supp } \mu, \{w_1, \dots, w_{n+1}\}) \leq \alpha_0\eta < 9 \cdot 2^{n+2}n^{2n+2}n^{15n} \varepsilon^{\frac{1}{4}} < n^{22n} \varepsilon^{\frac{1}{4}}. \quad (80)$$

Therefore we indirectly assume that

$$\zeta := \max_{u \in \text{supp } \mu} \min_{i=1, \dots, n+1} \angle(u, w_i) > \alpha_0\eta,$$

and hence there is some $u_0 \in \text{supp } \mu$ such that $\min_{i=1}^{n+1} \angle(u_0, w_i) = \zeta$. Let

$$L := \text{conv}\{u_0, u_1, \dots, u_{n+1}\}.$$

Lemma 5.5 and (75) imply

$$V(L^\circ) \leq \left(1 - \frac{\zeta}{2^{n+2}n^{2n}} \right) V(\Delta_n^\circ).$$

Since L is a polytope with $n + 2$ vertices, it is shown in Meyer, Reisner [57] that

$$V(L)V(L^\circ) \geq V(\Delta_n)V(\Delta_n^\circ),$$

which proves a special case of the Mahler conjecture. Therefore we get

$$V(L) \geq \left(1 + \frac{\zeta}{2^{n+2}n^{2n}} \right) V(\Delta_n), \quad (81)$$

while readily

$$\tilde{\Delta}_n \subset S_0 \subset L$$

holds for $\tilde{\Delta}_n = (1 + 2n\eta)^{-1}\Delta_n$. It follows from this, $L \subset B^n$ and (81) that

$$\begin{aligned} \gamma_n(tL) &> \gamma_n(t\tilde{\Delta}_n) \text{ for } t > 0, \\ \gamma_n(tL) &> \gamma_n(t\tilde{\Delta}_n) + \frac{e^{-\frac{1}{2}}}{\sqrt{2\pi^{\frac{n}{2}}}} \frac{t^n \zeta}{2^{n+2} n^{2n}} V(\Delta_n) \text{ for } t \in (0, 1]. \end{aligned}$$

We deduce from (2) and Lemma 5.7 (d) that

$$\ell(\tilde{\Delta}_n) - \ell(L) \geq \int_0^1 \gamma_n(tL) - \gamma_n(t\tilde{\Delta}_n) dt > \int_0^1 \frac{t^n \zeta}{n^{5n+5}} \ell(\Delta_n) dt > n^{-8n} \ell(\Delta_n) \zeta.$$

We conclude from (77) that

$$(1 - \varepsilon)\ell(\Delta_n) \leq \ell(Z_\infty(\mu)) \leq \ell(L) \leq (1 - n^{-8n}\zeta + 2n\eta)\ell(\Delta_n),$$

and hence

$$\zeta \leq n^{8n}(2n\eta + \varepsilon) < n^{22n}\varepsilon^{\frac{1}{4}}.$$

Therefore in both cases (compare (80)), if $\ell(Z_\infty(\mu)) \geq (1 - \varepsilon)\ell(\Delta_n)$ for a centered isotropic measure μ on S^{n-1} , then

$$\delta_H(\text{supp } \mu, \{w_1, \dots, w_{n+1}\}) < n^{28n}\varepsilon^{\frac{1}{4}}.$$

Since $\ell(Z_\infty(\mu)) \geq (1 - \varepsilon)\ell(\Delta_n)$ and $W(Z_\infty(\mu)^\circ) \geq (1 - \varepsilon)W(\Delta_n^\circ)$ are equivalent according to (1), we have verified the case $W(Z_\infty(\mu)^\circ) \geq (1 - \varepsilon)W(\Delta_n^\circ)$ of Theorem 1.6 as well, in the case $\varepsilon < n^{-100n}$. However, if $\varepsilon \geq n^{-100n}$, then Theorem 1.6 trivially holds since for any $x \in S^{n-1}$ there exists a vertex w of Δ_n with $\|x - w\| \leq \sqrt{2}$. \square

9 An almost regular simplex for Theorem 1.4 and Theorem 1.6 (c), (d)

The whole section is dedicated to proving the following statement.

Proposition 9.1 *Let $n + 1 \leq k \leq 2n^2$, $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ be such that*

$$\begin{aligned} \sum_{i=1}^k c_i u_i \otimes u_i &= I_n, \\ \sum_{i=1}^k c_i u_i &= o, \end{aligned}$$

and $\ell(C^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ holds for $C = \text{conv}\{u_1, \dots, u_k\}$ and $\varepsilon \in (0, n^{-60n})$.

Then for $\eta = n^{15n}\varepsilon^{\frac{1}{4}} \in (0, 1)$, there exists a regular simplex with vertices $w_1, \dots, w_{n+1} \in S^{n-1}$ and $\{i_1, \dots, i_{n+1}\} \subset \{1, \dots, k\}$ such that

$$\angle(u_{i_j}, w_j) \leq \eta \text{ for } j = 1, \dots, n + 1.$$

We recall from (66) that if $s \in \mathbb{R}$, then \tilde{g}_s is defined by

$$\tilde{g}_s(t) = \mathbf{1}\{t \geq 0\} \exp\left(-\frac{(t-s)^2}{2}\right), \quad t \in \mathbb{R}.$$

In this section, we slightly change the setup used in Proposition 6.2 and Corollary 6.3. As before Proposition 6.2, we fix an $e \in S^n \subset \mathbb{R}^{n+1}$, and identify $e^\perp \subset \mathbb{R}^{n+1}$ with \mathbb{R}^n . However now, for each $u_i \in S^{n-1}$, we consider

$$\begin{aligned} \tilde{u}_i &= -\frac{\sqrt{n}}{\sqrt{n+1}} u_i + \frac{1}{\sqrt{n+1}} e \in S^n, \\ \tilde{c}_i &= \frac{n+1}{n} c_i, \end{aligned}$$

and hence

$$\sum_{i=1}^k \tilde{c}_i \tilde{u}_i \otimes \tilde{u}_i = \mathbf{I}_{n+1}, \quad (82)$$

$$\sum_{i=1}^k \tilde{c}_i = n+1, \quad (83)$$

$$\sum_{i=1}^k \tilde{c}_i \tilde{u}_i = \frac{\sum_{i=1}^k \tilde{c}_i}{\sqrt{n+1}} \cdot e = \sqrt{n+1} e. \quad (84)$$

For the convex cone

$$\tilde{\mathcal{C}} := \{z \in \mathbb{R}^{n+1} : \langle z, \tilde{u}_i \rangle \geq 0 \ i = 1, \dots, k\},$$

the use of $-u_i$ instead of u_i in the definition of \tilde{u}_i ensures that

$$x + r e \in \tilde{\mathcal{C}} \text{ for } x \in \mathbb{R}^n \text{ and } r \in \mathbb{R} \text{ if and only if } r \geq 0 \text{ and } x \in \frac{r}{\sqrt{n}} C^\circ. \quad (85)$$

Moreover, we observe that if $C = \Delta_n$, then $k = n+1$, and $\tilde{u}_1, \dots, \tilde{u}_{n+1}$ form an orthonormal basis of \mathbb{R}^{n+1} .

Since $-u_1, \dots, -u_k$ satisfy the same conditions as u_1, \dots, u_k , it follows that Corollary 6.3 remains true for the vectors $\tilde{u}_1, \dots, \tilde{u}_k$ as defined in this section.

We suppose that Proposition 9.1 does not hold, and we seek a contradiction. From Corollary 6.3 and (67) we deduce that if $s \in [0, 0.15]$, then

$$\int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle z, \tilde{u}_i \rangle)^{\tilde{c}_i} dz \leq (1 - n^{-56n} \eta^4) \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} \leq \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} - n^{-56n} \eta^4. \quad (86)$$

Next we state a counterpart to Lemma 7.2, which provides general relations independent of the indirect reasoning used to establish Proposition 9.1.

Lemma 9.2 *If $s \in \mathbb{R}$ and C is defined as above, then*

$$(i) (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n \left(\frac{r}{\sqrt{n}} C^\circ \right) dr = \int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle z, \tilde{u}_i \rangle)^{\tilde{c}_i} dz,$$

$$(ii) (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n \left(\frac{r}{\sqrt{n}} \Delta_n^\circ \right) dr = \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1}.$$

Proof. Applying first (82), (83), (84) and then (85), we obtain

$$\begin{aligned} \int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle z, \tilde{u}_i \rangle)^{\tilde{c}_i} dz &= \int_{\tilde{c}} \exp \left(-\frac{1}{2} \sum_{i=1}^k \tilde{c}_i \langle z, \tilde{u}_i \rangle^2 + s \sum_{i=1}^k \tilde{c}_i \langle z, \tilde{u}_i \rangle - \frac{s^2}{2} \sum_{i=1}^k \tilde{c}_i \right) dz \\ &= e^{-\frac{(n+1)s^2}{2}} \int_{\tilde{c}} e^{-\frac{\|z\|^2}{2} + s\sqrt{n+1}\langle z, e \rangle} dz \\ &= e^{-\frac{(n+1)s^2}{2}} \int_0^\infty \int_{\frac{r}{\sqrt{n}} C^\circ} e^{-\frac{\|x\|^2 + r^2}{2} + sr\sqrt{n+1}} dx dr \\ &= (2\pi)^{\frac{n}{2}} e^{-\frac{(n+1)s^2}{2}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n \left(\frac{r}{\sqrt{n}} C^\circ \right) dr. \end{aligned}$$

For (ii), we observe that if we replace C by Δ_n in the argument above, then the analogues of $\tilde{u}_1, \dots, \tilde{u}_{n+1}$ form an orthonormal basis of \mathbb{R}^{n+1} and \tilde{c}_i is replaced by 1. \square

We apply the change of parameter $\tau = s\sqrt{\frac{n+1}{n}}$ and the substitution $t = r/\sqrt{n}$ in Lemma 9.2, and conclude with the help of the Brascamp-Lieb inequality (13) that if $\tau \in \mathbb{R}$, then

$$\begin{aligned} \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} \gamma_n(tC^\circ) dt &= e^{-\frac{n\tau^2}{2}} \int_0^\infty e^{-\frac{nt^2}{2} + st\sqrt{n(n+1)}} \gamma_n(tC^\circ) dt \\ &= \frac{e^{-\frac{(n+1)s^2}{2}}}{\sqrt{n}} \int_0^\infty e^{-\frac{r^2}{2} + sr\sqrt{n+1}} \gamma_n \left(\frac{r}{\sqrt{n}} C^\circ \right) dr \\ &= \frac{1}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle z, \tilde{u}_i \rangle)^{\tilde{c}_i} dz \\ &\leq \frac{1}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} \\ &= \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} \gamma_n(t\Delta_n^\circ) dt, \end{aligned}$$

and hence

$$\int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(t\Delta_n^\circ)) dt \leq \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(tC^\circ)) dt. \quad (87)$$

In addition, if $\tau \in [0, 0.15] \subset [0, 0.15\sqrt{\frac{n+1}{n}}]$, which implies that $s \in [0, 0.15]$, then using (86) instead of the Brascamp-Lieb inequality (13), we obtain

$$\begin{aligned} \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} \gamma_n(tC^\circ) dt &= \frac{1}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^{n+1}} \prod_{i=1}^k \tilde{g}_s(\langle z, \tilde{u}_i \rangle)^{\tilde{c}_i} dz \\ &\leq \frac{1}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \left(\left(\int_{\mathbb{R}} \tilde{g}_s \right)^{n+1} - n^{-56n} \eta^4 \right) \\ &= \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} \gamma_n(t\Delta_n^\circ) dt - \frac{n^{-56n}}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \eta^4. \end{aligned}$$

Hence, we get

$$\int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(t\Delta_n^\circ)) dt \leq \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(tC^\circ)) dt - \frac{0.15 n^{-56n}}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \eta^4. \quad (88)$$

Integrating (87) for $\tau \in \mathbb{R} \setminus [0, 0.15]$ and (88) for $\tau \in [0, 0.15]$, we deduce that

$$\begin{aligned} \int_{-\infty}^\infty \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(t\Delta_n^\circ)) dt d\tau \\ \leq \int_{-\infty}^\infty \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(tC^\circ)) dt d\tau - \frac{0.15 n^{-56n}}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \eta^4. \end{aligned} \quad (89)$$

Since for any $t \in \mathbb{R}$, we have

$$\int_{-\infty}^\infty e^{-\frac{n}{2}(t-\tau)^2} d\tau = \sqrt{\frac{2\pi}{n}},$$

we deduce from (2) and (89) that

$$\begin{aligned} \ell(C^\circ) &= \int_0^\infty (1 - \gamma_n(tC^\circ)) dt \\ &= \sqrt{\frac{n}{2\pi}} \int_{-\infty}^\infty \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(tC^\circ)) dt d\tau \\ &\geq \sqrt{\frac{n}{2\pi}} \int_{-\infty}^\infty \int_0^\infty e^{-\frac{n}{2}(t-\tau)^2} (1 - \gamma_n(t\Delta_n^\circ)) dt d\tau + \frac{0.15 n^{-56n}}{\sqrt{n}(2\pi)^{\frac{n}{2}}} \eta^4 \\ &= \int_0^\infty (1 - \gamma_n(t\Delta_n^\circ)) dt + \frac{0.15 n^{-56n}}{(2\pi)^{\frac{n+1}{2}}} \eta^4 \\ &\geq \ell(\Delta_n^\circ) + n^{-60n} \eta^4 \ell(\Delta_n^\circ), \end{aligned}$$

where Lemma 5.7 (a) was used in the last step. This shows that

$$(1 + \varepsilon) \ell(\Delta_n^\circ) \geq \ell(C^\circ) > (1 + n^{-60n} \eta^4) \ell(\Delta_n^\circ),$$

which contradicts $\eta = n^{15n} \varepsilon^{\frac{1}{4}}$, and in turn implies Proposition 9.1.

10 Proof of Theorem 1.4 and of Theorem 1.6 (c), (d)

For Theorem 1.6, let μ be a centered isotropic measure on S^{n-1} , and let $K = Z_\infty(\mu)^\circ$, and hence $\text{supp } \mu = \partial K \cap S^{n-1}$. In particular, under the assumptions of Theorem 1.4 and of Theorem 1.6 (d), we have $\ell(K) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$. First, we assume that

$$\varepsilon < n^{-100n}.$$

It follows from Lemma 2.1 and John's theorem that there exist $k \geq n + 1$ with $k \leq 2n^2$, $u_1, \dots, u_k \in \partial K \cap S^{n-1}$ and $c_1, \dots, c_k > 0$ such that

$$\begin{aligned} \sum_{i=1}^k c_i u_i \otimes u_i &= I_n, \\ \sum_{i=1}^k c_i u_i &= o. \end{aligned}$$

We write μ_0 to denote the centered discrete isotropic measure with $\text{supp } \mu_0 = \{u_1, \dots, u_k\}$ and $\mu_0(\{u_i\}) = c_i$ for $i = 1, \dots, k$, and define (again)

$$C := Z_\infty(\mu_0) = \text{conv}\{u_1, \dots, u_k\} \subset K^\circ.$$

Since $\ell(C^\circ) \leq \ell(K) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$, it follows from Proposition 9.1 that we may assume that the vertices w_1, \dots, w_{n+1} of Δ_n satisfy

$$\angle(u_i, w_i) \leq \eta \text{ for } \eta = n^{15n} \varepsilon^{\frac{1}{4}} \quad \text{and} \quad i = 1, \dots, n + 1. \quad (90)$$

We observe that $K \subset S_1 := S_0^\circ$, where S_1 is the polar of S_0 and the facets of

$$S_1 = \bigcap_{i=1}^{n+1} \{x \in \mathbb{R}^n : \langle x, u_i \rangle \leq 1\}$$

touch B^n at u_1, \dots, u_{n+1} . We deduce from (90) and Lemma 5.4 that

$$(1 - n\eta)\Delta_n^\circ \subset S_1 \subset (1 + 2n\eta)\Delta_n^\circ \subset 2\Delta_n^\circ. \quad (91)$$

We claim that

$$\delta_{\text{vol}}(K, S_1) = V(S_1 \setminus K) \leq n^{23n} \varepsilon^{\frac{1}{4}}. \quad (92)$$

Using $\frac{1}{2n} S_1 \subset \frac{1}{n} \Delta_n^\circ \subset B^n$, (2) and Lemma 5.7 (a), we get

$$\begin{aligned} \ell(K) - \ell(S_1) &= \int_0^\infty (\gamma_n(tS_1) - \gamma_n(tK)) dt \geq \int_0^{\frac{1}{2n}} \frac{e^{-\frac{1}{2}}}{(2\pi)^{n/2}} t^n V(S_1 \setminus K) dt \\ &\geq \frac{e^{-\frac{1}{2}}}{(n+1)(2\pi)^{n/2}(2n)^{n+1}} V(S_1 \setminus K) \frac{\ell(\Delta_n^\circ)}{\sqrt{n}} > \frac{V(S_1 \setminus K)}{n^{6n}} \ell(\Delta_n^\circ). \end{aligned} \quad (93)$$

In addition, (91) yields

$$\ell(S_1) - \ell(\Delta_n^\circ) \geq \ell((1 + 2n\eta)\Delta_n^\circ) - \ell(\Delta_n^\circ) = ((1 + 2n\eta)^{-1} - 1)\ell(\Delta_n^\circ) \geq -2n\eta \ell(\Delta_n^\circ). \quad (94)$$

We deduce from $\ell(K) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$, (93) and (94) that

$$\varepsilon \ell(\Delta_n^\circ) \geq \ell(K) - \ell(\Delta_n^\circ) > \left(\frac{V(S_1 \setminus K)}{n^{6n}} - 2n\eta \right) \ell(\Delta_n^\circ).$$

Then $\eta = n^{15n}\varepsilon^{\frac{1}{4}}$ implies that

$$V(S_1 \setminus K) < n^{6n}(\varepsilon + 2n\eta) < 4n \cdot n^{6n}\eta < n^{23n}\varepsilon^{\frac{1}{4}},$$

which proves (92).

Proof of Theorem 1.4: We start to deal with the symmetric volume distance of K and Δ° .

Using (91), $(1 + 2n\eta)^n \leq 4/3$ and Lemma 5.7 (b), we get

$$\delta_{\text{vol}}(S_0, \Delta_n^\circ) \leq ((1 + 2n\eta)^n - (1 - n\eta)^n) V(\Delta_n^\circ) < n \cdot 4n\eta V(\Delta_n^\circ) < n^{19n}\varepsilon^{\frac{1}{4}}. \quad (95)$$

Combining (92) and (95), we get $\delta_{\text{vol}}(K, \Delta^\circ) \leq n^{24n}\varepsilon^{\frac{1}{4}}$, which proves Theorem 1.4 (i) under the assumption $\varepsilon < n^{-100n}$.

In order to derive an upper bound for the Hausdorff distance of K and Δ° , we first show that the centroid σ_0 of S_0 satisfies

$$\sigma_0 \in 4n\eta\Delta_n^\circ. \quad (96)$$

To prove (96), we observe that

$$\Delta_n^\circ = \bigcap_{i=1}^{n+1} \{x \in \mathbb{R}^n : \langle x, w_i \rangle \leq 1\} = \text{conv}\{-nw_1, \dots, -nw_{n+1}\}.$$

For each $j \in \{1, \dots, n+1\}$, (91) yields that S_1 has a vertex v_j with

$$\langle -w_j, v_j \rangle \geq h_{(1-n\eta)\Delta_n^\circ}(-w_j) = (1 - n\eta)n.$$

Since $S_1 \subset (1 + 2n\eta)\Delta_n^\circ$ and $\Delta_n^\circ + nw_j$ is homothetic to Δ_n° with o as the vertex with exterior normal $-w_j$, we have

$$\begin{aligned} v_j &\in \{x \in (1 + 2n\eta)\Delta_n^\circ : \langle -w_j, x \rangle \geq (1 - n\eta)n\} \\ &= -(1 + 2n\eta)nw_j + 3n\eta((1 + 2n\eta)\Delta_n^\circ + (1 + 2n\eta)nw_j) \end{aligned} \quad (97)$$

for $j = 1, \dots, n+1$. Hence, the vertices v_1, \dots, v_{n+1} are contained in mutually disjoint neighbourhoods of $-nw_1, \dots, -nw_{n+1}$ and thus $S_1 = \text{conv}\{v_1, \dots, v_{n+1}\}$.

If $i = 1, \dots, n+1$, then $\langle w_i, w_j \rangle = \frac{-1}{n}$ for $j \neq i$ implies $\langle w_i, x \rangle \leq n+1$ for $x \in \Delta_n^\circ + nw_i$ and $\langle w_i, y \rangle \leq 0$ for $y \in \Delta_n^\circ + nw_j$ and $j \neq i$. Therefore (97) yields

$$(n+1)\langle w_i, \sigma_0 \rangle = \langle w_i, v_i \rangle + \sum_{j \neq i} \langle w_i, v_j \rangle \leq (1 + 2n\eta)[-n + 3n\eta(n+1)] + n(1 + 2n\eta)$$

$$\leq 3n(n+1)(1+2n\eta)\eta \leq 4n(n+1)\eta,$$

for $i = 1, \dots, n+1$, which proves the claim.

Note that $\sigma_0 \in 4n\eta\Delta_n^\circ \subset 4n^2\eta B^n \subset \text{int}(B^n) \subset K \subset S_1$, in particular we have $o \in \text{int}(K - \sigma_0)$ and $K - \sigma_0 \subset S_1 - \sigma_0$. Let $\xi \in [0, 1)$ be minimal such that

$$\sigma_0 + (1 - \xi)(S_1 - \sigma_0) \subset K.$$

From (91), $\eta < 1/(4n^2)$ and Lemma 5.7 (c), we deduce that

$$V(S_1) \geq (1 - n\eta)^n n^n V(\Delta_n) \geq \left[\left(1 - \frac{1}{4n}\right)^2 \left(1 + \frac{1}{n}\right) \right]^{\frac{n}{2}} > 1.$$

Then it follows from Lemma 5.6 (i) and (92) that

$$\frac{\xi^n}{e} V(S_1) \leq V(S_1 \setminus K) < n^{23n} \varepsilon^{\frac{1}{4}} < n^{23n} \varepsilon^{\frac{1}{4}} V(S_1),$$

and hence $\xi < n^{24} \varepsilon^{\frac{1}{4n}}$. We deduce from (96) that $-\sigma_0 \in 4n^2\eta\Delta_n^\circ$, and therefore

$$S_1 - \sigma_0 \subset (1 + 2n\eta)\Delta_n^\circ + 4n^2\eta\Delta_n^\circ \subset 2\Delta_n^\circ \subset 2nB^n.$$

Since $K \subset S_1 \subset K + \xi(S_1 - \sigma_0)$ by the definition of ξ , it follows that

$$\delta_H(S_1, K) \leq 2n\xi < n^{26} \varepsilon^{\frac{1}{4n}}.$$

On the other hand, $\eta = n^{15n} \varepsilon^{\frac{1}{4}}$, (91) and $\Delta_n^\circ \subset nB^n$ imply that

$$\delta_H(S_1, \Delta_n^\circ) < 2n^2\eta < n^{17n} \varepsilon^{\frac{1}{4}}.$$

Since $n^{17n} \varepsilon^{\frac{1}{4}} < n^{26} \varepsilon^{\frac{1}{4n}}$ if $\varepsilon < n^{-100n}$, we have $\delta_H(K, \Delta_n^\circ) < n^{27} \varepsilon^{\frac{1}{4n}}$, which completes the proof of Theorem 1.4 if $\varepsilon < n^{-100n}$.

However, if $\varepsilon \geq n^{-100n}$, then Theorem 1.4 trivially holds as $\delta_{\text{vol}}(M, \Delta_n) \leq n^n \kappa_n$ and $\delta_H(M, \Delta_n) \leq n$ for any convex body $M \subset nB^n$. Note that if B^n is the John ellipsoid of K , then $K \subset nB^n$, and $\kappa_n \leq 6$ for all $n \in \mathbb{N}$. \square

Proof of Theorem 1.6 (c), (d): Suppose that $\ell(Z_\infty(\mu)^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$.

Let $\alpha_0 = 9 \cdot 2^{n+2} n^{2n+2}$ be the constant of Lemma 5.5. If for any $u \in \text{supp } \mu$ there exists a w_i such that $\angle(u, w_i) \leq \alpha_0 \eta$, then

$$\delta_H(\text{supp } \mu, \{w_1, \dots, w_{n+1}\}) \leq \alpha_0 \eta < 9 \cdot 2^{n+2} n^{2n+2} n^{15n} \varepsilon^{\frac{1}{4}} < n^{22n} \varepsilon^{\frac{1}{4}}.$$

Therefore we assume that

$$\zeta := \max_{u \in \text{supp } \mu} \min_{i=1, \dots, n+1} \angle(u, w_i) > \alpha_0 \eta,$$

and let $u_0 \in \text{supp } \mu$ be such that $\min\{\angle(u_0, w_i) : i = 1, \dots, n+1\} = \zeta$. Let

$$L = \text{conv}\{u_0, u_1, \dots, u_{n+1}\},$$

and hence $Z_\infty(\mu)^\circ = K \subset L^\circ \subset S_1$. Lemma 5.5 and (90) imply

$$V(L^\circ) \leq \left(1 - \frac{\zeta}{2^{n+2}n^{2n}}\right) V(\Delta_n^\circ),$$

thus

$$\delta_{\text{vol}}(L^\circ, \Delta_n^\circ) \geq \frac{\zeta}{2^{n+2}n^{2n}} V(\Delta_n^\circ).$$

On the other hand, (91) and $(1 + 2n\eta)^n < 4/3$ yield

$$\delta_{\text{vol}}(S_1, \Delta_n^\circ) \leq ((1 + 2n\eta)^n - (1 - n\eta)^n) V(\Delta_n^\circ) < 4n^2\eta V(\Delta_n^\circ).$$

Therefore the triangle inequality implies that

$$V(S_1 \setminus L^\circ) = \delta_{\text{vol}}(S_1, L^\circ) \geq \left(\frac{\zeta}{2^{n+2}n^{2n}} - 4n^2\eta\right) V(\Delta_n^\circ).$$

Since $V(\Delta_n^\circ) > 1$ by Lemma 5.7 (c), we deduce from (92) that

$$n^{23n}\varepsilon^{\frac{1}{4}} V(\Delta_n^\circ) \geq V(S_1 \setminus Z_\infty(\mu)^\circ) \geq V(S_1 \setminus L^\circ) > \left(\frac{\zeta}{2^{n+2}n^{2n}} - 4n^2\eta\right) V(\Delta_n^\circ).$$

It follows from $\eta = n^{15n}\varepsilon^{\frac{1}{4}}$ that

$$\zeta < 2^{n+2}n^{2n}(n^{23n}\varepsilon^{\frac{1}{4}} + 4n^2\eta) < n^{28n}\varepsilon^{\frac{1}{4}},$$

which proves Theorem 1.6 in the case where $\ell(Z_\infty(\mu)^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ and $\varepsilon < n^{-100n}$.

Since $\ell(Z_\infty(\mu)^\circ) \leq (1 + \varepsilon)\ell(\Delta_n^\circ)$ and $W(Z_\infty(\mu)) \leq (1 + \varepsilon)W(\Delta_n)$ are equivalent according to (1), we have completed the proof of Theorem 1.6 if $\varepsilon < n^{-100n}$.

However, if $\varepsilon \geq n^{-100n}$, then Theorem 1.6 trivially holds as for any $x \in S^{n-1}$ there exists a vertex w of Δ_n with $\|x - w\| \leq \sqrt{2}$. \square

11 Proof of Corollary 1.5

For the proof of Corollary 1.5, we need the following observation.

Lemma 11.1 *If $\frac{1}{n}B^n \subset K, C \subset nB^n$ for convex bodies K and C in \mathbb{R}^n , then*

$$\frac{1}{n^2} \delta_H(K, C) \leq \delta_H(K^\circ, C^\circ) \leq n^2 \delta_H(K, C).$$

Proof. We also have $\frac{1}{n} B^n \subset K^\circ, C^\circ \subset nB^n$. First, we show

$$\delta_H(K^\circ, C^\circ) \leq n^2 \delta_H(K, C). \quad (98)$$

Since $K \subset C + \delta_H(K, C)B^n \subset C + n\delta_H(K, C)C = (1 + n\delta_H(K, C))C$, we have

$$C^\circ \subset (1 + n\delta_H(K, C))K^\circ \subset K^\circ + n^2\delta_H(K, C)B^n.$$

By symmetry, we also have $K^\circ \subset C^\circ + n^2\delta_H(K, C)B^n$, and thus we have verified (98).

Changing the roles of K, C and their polars K°, C° in (98) (and using the bipolar theorem), we also deduce the inequality $\delta_H(K, C) \leq n^2\delta_H(K^\circ, C^\circ)$. \square

Since $W(K) = \frac{2}{\ell(B^n)} \ell(K^\circ)$ according to (1), we conclude Corollary 1.5 by combining Theorem 1.3 (ii), Theorem 1.4 (ii) and Lemma 11.1. \square

Remark The factor n^2 in Lemma 11.1 is optimal.

References

- [1] D. Alonso-Gutiérrez, J. Bastero: Approaching the Kannan-Lovász-Simonovits and variance conjectures. Lecture Notes in Mathematics, 2131. Springer, Cham, 2015.
- [2] S. Artstein-Avidan, A. Giannopoulos, V.D. Milman: Asymptotic geometric analysis. Part I. Mathematical Surveys and Monographs, 202. American Mathematical Society, Providence, RI, 2015.
- [3] K. Ball: Volumes of sections of cubes and related problems. In: J. Lindenstrauss and V.D. Milman (ed), Israel seminar on Geometric Aspects of Functional Analysis (1987–1988), 1376, Lectures Notes in Mathematics. Springer-Verlag, 1989.
- [4] K. Ball: Volume ratios and a reverse isoperimetric inequality. J. London Math. Soc. 44 (1991), 351–359.
- [5] K. Ball: Shadows of convex bodies. Trans. Amer. Math. Soc. 327 (1991), 891–901.
- [6] K. Ball: Ellipsoids of maximal volume in convex bodies. Geom. Dedicata 41 (1992), 241–250.
- [7] K. Ball: Convex geometry and functional analysis. In: W.B. Johnson, L. Lindenstrauss (eds), Handbook of the geometry of Banach spaces, 1, (2003), 161–194.
- [8] Z. Balogh, A. Kristály: Equality in Borell-Brascamp-Lieb inequalities on curved spaces. Adv. Math. 339 (2018), 453–494.
- [9] F. Barthe: Inégalités de Brascamp-Lieb et convexité. C. R. Acad. Sci. Paris 324 (1997), 885–888.

- [10] F. Barthe: On a reverse form of the Brascamp-Lieb inequality. *Invent. Math.* 134 (1998), 335–361.
- [11] F. Barthe: An extremal property of the mean width of the simplex. *Math. Ann.* 310 (1998), 685–693.
- [12] F. Barthe: A continuous version of the Brascamp-Lieb inequalities. *Geometric Aspects of Functional Analysis, Lecture Notes in Mathematics Volume, 1850, 2004*, 53–63.
- [13] F. Barthe, D. Cordero-Erausquin: Invariances in variance estimates. *Proc. Lond. Math. Soc.* 106 (2013), 33–64.
- [14] F. Barthe, D. Cordero-Erausquin, M. Ledoux, B. Maurey: Correlation and Brascamp-Lieb inequalities for Markov semigroups. *Int. Math. Res. Not.* 10 (2011), 2177–2216.
- [15] F. Behrend: Über einige Affinvarianten konvexer Bereiche. (German) *Math. Ann.* 113 (1937), 713–747.
- [16] J. Bennett, N. Bez, T.C. Flock, S. Lee: Stability of the Brascamp-Lieb constant and applications. *Amer. J. Math.* 140 (2018), 543–569.
- [17] J. Bennett, T. Carbery, M. Christ, T. Tao: The Brascamp–Lieb Inequalities: Finiteness, Structure and Extremals. *Geom. Func. Anal.* 17 (2008), 1343–1415.
- [18] C. Borell: Convex set functions in d -space. *Period. Math. Hung.* 6 (1975), no. 2, 111–136.
- [19] K. Böröczky, Jr.: Some extremal properties of the regular simplex. *Intuitive geometry. Colloq. Math. Soc. János Bolyai, 63, North-Holland, Amsterdam, (1994)*, 45–61.
- [20] K.J. Böröczky, D. Hug: Isotropic measures and stronger forms of the reverse isoperimetric inequality. *Trans. Amer. Math. Soc.* 369 (2017), 6987–7019.
- [21] K.J. Böröczky, E. Lutwak, D. Yang, G. Zhang: Affine images of isotropic measures. *J. Diff. Geom.* 99 (2015), 407–442.
- [22] H.J. Brascamp, E.H. Lieb: Best constants in Young’s inequality, its converse, and its generalization to more than three functions. *Advances in Math.* 20 (1976), 151–173.
- [23] E. Carlen, D. Cordero-Erausquin: Subadditivity of the entropy and its relation to Brascamp-Lieb type inequalities. *Geom. Funct. Anal.* 19 (2009), 373–405.
- [24] L. Dalla, D.G. Larman, P. Mani-Levitska, C. Zong: The blocking numbers of convex bodies. *Discrete Comput. Geom.* 24 (2002), 267–277.
- [25] V.I. Diskant: Stability of the solution of a Minkowski equation. (Russian) *Sibirsk. Mat. Ž.* 14 (1973), 669–673. [Eng. transl.: *Siberian Math. J.* 14 (1974), 466–473.]
- [26] L. Dümbgen: Bounding Standard Gaussian Tail Probabilities. [arxiv:1012.2063v3](https://arxiv.org/abs/1012.2063v3)

- [27] L. Fejes Tóth: Lagerungen in der Ebene, auf der Kugel und im Raum. Springer-Verlag, Berlin, 2nd edition, 1972.
- [28] A. Figalli, F. Maggi, A. Pratelli: A refined Brunn-Minkowski inequality for convex sets. *Annales de IHP (C) Non Linear Analysis* 26 (2009), 2511–2519.
- [29] A. Figalli, F. Maggi, A. Pratelli: A mass transportation approach to quantitative isoperimetric inequalities. *Inventiones Mathematicae* 182, (2010), 167–211.
- [30] B. Fleury, O. Guédon, G. Paouris: A stability result for mean width of L_p -centroid bodies. *Adv. Math.* 214 (2007), no. 2, 865–877.
- [31] N. Fusco, F. Maggi, A. Pratelli: The sharp quantitative isoperimetric inequality. *Ann. of Math.* 168 (2008), no. 3, 941–980.
- [32] D. Ghilli, P. Salani: Quantitative Borell-Brascamp-Lieb inequalities for power concave functions. *J. Convex Anal.* 24 (2017), 857–888.
- [33] A.A. Giannopoulos, V.D. Milman, M. Rudelson: Convex bodies with minimal mean width. In: *Geometric aspects of functional analysis, Lecture Notes in Math.*, 1745, Springer, Berlin, 2000, 81–93.
- [34] A.A. Giannopoulos, M. Papadimitrakis: Isotropic surface area measures. *Mathematika* 46 (1999), 1–13
- [35] R.D. Gordon: Values of Mills’ ratio of area to bounding ordinate and of the normal probability integral for large values of the argument. *Annals of Mathematical Statistics* 12 (1941), 364–366.
- [36] H. Groemer: Stability properties of geometric inequalities. *Amer. Math. Monthly* 97 (1990), no. 5, 382–394.
- [37] H. Groemer: Stability of geometric inequalities. *Handbook of convex geometry, Vol. A, B*, 125–150, North-Holland, Amsterdam, 1993.
- [38] H. Groemer, R. Schneider: Stability estimates for some geometric inequalities. *Bull. London Math. Soc.* 23 (1991), no. 1, 67–74.
- [39] P.M. Gruber: *Convex and discrete geometry. Grundlehren der Mathematischen Wissenschaften*, Springer, Berlin, 2007.
- [40] P.M. Gruber, F.E. Schuster: An arithmetic proof of John’s ellipsoid theorem. *Arch. Math.* 85 (2005), 82–88.
- [41] B. Grünbaum: Partitions of mass-distributions and of convex bodies by hyperplanes, *Pacific J. Math.* 10 (1960), 1257–1261.

- [42] O. Guedon, E. Milman: Interpolating thin-shell and sharp large-deviation estimates for isotropic log-concave measures. *Geom. Funct. Anal.* 21 (2011), 1043–1068.
- [43] D. Hug, R. Schneider: Reverse inequalities for zonoids and their application. *Adv. Math.* 228 (2011), 2634–2646.
- [44] F. John: Polar correspondence with respect to a convex region. *Duke Math. J.* 3 (1937), 355–369.
- [45] F. John: Extremum problems with inequalities as subsidiary conditions. In: *Studies and Essays Presented to R. Courant on His 60th Birthday, January 8, 1948*, pp. 187–204, Interscience Publishers, New York, 1948.
- [46] R. Kannan, L. Lovász, M. Simonovits: Isoperimetric problems for convex bodies and a localization lemma. *Discrete Comput. Geom.* 13 (1995), 541–559.
- [47] B. Klartag: A Berry-Esseen type inequality for convex bodies with an unconditional basis, *Probab. Theory Related Fields* 145 (2009), 1–33.
- [48] B. Klartag: On nearly radial marginals of high-dimensional probability measures, *J. Eur. Math. Soc.*, 12 (2010), 723–754.
- [49] A.-J. Li, G. Leng: Mean width inequalities for isotropic measures. *Math. Z.* 270 (2012), 1089–1110.
- [50] E.H. Lieb: Gaussian kernels have only Gaussian maximizers. *Invent. Math.* 102 (1990), 179–208.
- [51] A.E. Litvak: Around the simplex mean width conjecture. In: *Analytic aspects of convexity*, Springer INdAM Ser. 25, (2018), 73–84.
- [52] E. Lutwak: Selected affine isoperimetric inequalities. In: *Handbook of convex geometry*, North-Holland, Amsterdam, 1993, 151–176.
- [53] E. Lutwak, D. Yang, G. Zhang: Volume inequalities for subspaces of L_p . *J. Diff. Geom.* 68 (2004), 159–184.
- [54] E. Lutwak, D. Yang, G. Zhang: L_p John ellipsoids. *Proc. London Math. Soc.* 90 (2005), 497–520.
- [55] E. Lutwak, D. Yang, G. Zhang: Volume inequalities for isotropic measures. *Amer. J. Math.* 129 (2007), 1711–1723.
- [56] T. Ma: The characteristic properties of the minimal L_p -mean width. *J. Funct. Spaces* (2017), Art. ID 2943073, 10 pp.
- [57] M. Meyer, S. Reisner: Shadow systems and volumes of polar convex bodies. *Mathematika* 53 (2006), 129–148.

- [58] E. Milman: On the mean-width of isotropic convex bodies and their associated L_p -centroid bodies. *Int. Math. Res. Not. IMRN* (2015), no. 11, 3408–3423.
- [59] C. M. Petty: Surface area of a convex body under affine transformations. *Proc. Amer. Math. Soc.* 12 (1961), 824–828.
- [60] A. Rossi, P. Salani: Stability for Borell-Brascamp-Lieb inequalities. *Geometric aspects of functional analysis*, Lecture Notes in Math., 2169, Springer, Cham, (2017), 339–363.
- [61] G. Schechtman, M. Schmuckenschläger: A concentration inequality for harmonic measures. In: *Geometric Aspects of Functional Analysis (1992-1994)*, Birkhauser, (1995), 255–273.
- [62] M. Schmuckenschläger: An extremal property of the regular simplex. In: *Convex geometric analysis*. Cambridge, (1999), 199–202.
- [63] R. Schneider: *Convex bodies: the Brunn-Minkowski theory. Second expanded edition. Encyclopedia of Mathematics and its Applications*, 151. Cambridge University Press, Cambridge, 2014.
- [64] G.C. Shephard, R.J. Webster: *Metrics for sets of convex bodies. Mathematika* 12 (1965), 73–88.
- [65] M. Weberndorfer: Shadow systems of asymmetric L_p zonotopes. *Adv. Math.* 240 (2013), 613–635.
- [66] J. G. Wendel, Note on the Gamma function, *Amer. Math. Monthly* 55 (9) (1948), 563—564.

Authors' addresses:

Károly J. Böröczky, MTA Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, Reáltanoda u. 13-15, 1053 Budapest, Hungary. E-mail: carlos@renyi.hu

Ferenc Fodor, Department of Geometry, Bolyai Institute, University of Szeged, Aradi vértanúk tere 1, 6720 Szeged, Hungary. E-mail: fodorf@math.u-szeged.hu

Daniel Hug, Karlsruhe Institute of Technology (KIT), D-76128 Karlsruhe, Germany. E-mail: daniel.hug@kit.edu