



The Planktonic Core Microbiome and Core Functions in the Cattle Rumen by Next Generation Sequencing

Roland Wirth¹, Gyula Kádár², Balázs Kakuk¹, Gergely Maróti³, Zoltán Bagi¹, Árpád Szilágyi¹, Gábor Rákhelyi^{1,4}, József Horváth⁵ and Kornél L. Kovács^{1,4,6*}

¹ Department of Biotechnology, University of Szeged, Szeged, Hungary, ² Helvecia Protein Ltd., Szeghalom, Hungary, ³ Institute of Plant Biology, Biological Research Center, Hungarian Academy of Sciences, Szeged, Hungary, ⁴ Institute of Biophysics, Biological Research Center, Hungarian Academy of Sciences, Szeged, Hungary, ⁵ Faculty of Agriculture, University of Szeged, Hódmezővásárhely, Hungary, ⁶ Department of Oral Biology and Experimental Dental Research, University of Szeged, Szeged, Hungary

OPEN ACCESS

Edited by:

Roland Hatzenpichler,
Montana State University,
United States

Reviewed by:

Mick Watson,
The University of Edinburgh,
United Kingdom
Stephan Schmitz-Esser,
Iowa State University, United States

*Correspondence:

Kornél L. Kovács
kornel.kovacs@bio.u-szeged.hu

Specialty section:

This article was submitted to
Microbial Physiology and Metabolism,
a section of the journal
Frontiers in Microbiology

Received: 10 April 2018

Accepted: 07 September 2018

Published: 24 September 2018

Citation:

Wirth R, Kádár G, Kakuk B, Maróti G,
Bagi Z, Szilágyi Á, Rákhelyi G,
Horváth J and Kovács KL (2018) The
Planktonic Core Microbiome
and Core Functions in the Cattle
Rumen by Next Generation
Sequencing. *Front. Microbiol.* 9:2285.
doi: 10.3389/fmicb.2018.02285

The cow rumen harbors a great variety of diverse microbes, which form a complex, organized community. Understanding the behavior of this multifarious network is crucial in improving ruminant nutrient use efficiency. The aim of this study was to expand our knowledge by examining 10 Holstein dairy cow rumen fluid fraction whole metagenome and transcriptome datasets. DNA and mRNA sequence data, generated by Ion Torrent, was subjected to quality control and filtering before analysis for core elements. The taxonomic core microbiome consisted of 48 genera belonging to *Bacteria* (47) and *Archaea* (1). The genus *Prevotella* predominated the planktonic core community. Core functional groups were identified using co-occurrence analysis and resulted in 587 genes, from which 62 could be assigned to metabolic functions. Although this was a minimal functional core, it revealed key enzymes participating in various metabolic processes. A diverse and rich collection of enzymes involved in carbohydrate metabolism and other functions were identified. Transcripts coding for enzymes active in methanogenesis made up 1% of the core functions. The genera associated with the core enzyme functions were also identified. Linking genera to functions showed that the main metabolic pathways are primarily provided by *Bacteria* and several genera may serve as a “back-up” team for the central functions. The key actors in most essential metabolic routes belong to the genus *Prevotella*. Confirming earlier studies, the genus *Methanobrevibacter* carries out the overwhelming majority of rumen methanogenesis and therefore methane emission mitigation seems conceivable via targeting the hydrogenotrophic methanogenesis.

Keywords: rumen, whole genome, whole transcriptome, core microbiome, core functions, metabolic pathways, functions, taxa

INTRODUCTION

Ruminants are one of the most successful groups of herbivorous mammals on the planet, having evolved the forestomach, the rumen, where feed is degraded before it enters the true stomach and the rest of the digestive system. The rumen provides an environment for a diverse consortium of anaerobic microbes. These microbes produce enzymes which are needed to break

down complex molecules, primarily plant polysaccharides (Dai et al., 2015). The decomposition of cellulose-rich fibers involves intimate symbiotic relationships in the microbiota (Moran, 2005). The hydrolysis of lignocellulose often limits the kinetics and efficiency of metabolite production (Chapleur et al., 2014; Güllert et al., 2016). During rumination, sophisticated mechanical pretreatment of the lignocellulosic substrate and the delicate oxygen gradient along the route of feed play important roles in making the degradation of recalcitrant substances more efficient (Bayané and Guiot, 2011). The rumen microbiome is a well-studied, although not thoroughly understood microbial ecosystem, which is able to utilize the plant material in an organized food chain (Krause et al., 2003; Chapleur et al., 2014). In the first step, *Protozoa*, *Fungi*, and *Bacteria* carry out the hydrolysis of polymers, such as cellulose and other complex carbohydrates, proteins and lipids to low molecular weight compounds (Lynd et al., 2002; Jindou et al., 2008; Güllert et al., 2016). Subsequently, fermentative bacteria convert these metabolites to short chain fatty acids (SCFAs) like acetate, propionate and butyrate, CO₂, H₂, alcohols and other compounds. Methanogenic *Archaea* produce methane in the final step of the microbial food-chain (Le Van et al., 1998; Hegarty and Gerdes, 1999; Lopez et al., 1999; Janssen and Kirs, 2008; Hook et al., 2010). These communities co-evolve with their host according to their ability to convert some of the SCFAs, methylamines, CO₂ and H₂ to CH₄, CO₂ and water (Jami and Mizrahi, 2012; Meale et al., 2012; Poulsen et al., 2013; Jewell et al., 2015). The host benefits most if the process generates SCFAs (Weimer, 2015), which are absorbed through the rumen wall and used in the biosynthesis of sugars, lipids, and amino acids for the animal. Nonetheless, the thermodynamic and redox balances function properly only if a certain amount of chemical energy is released in the form of biogas, i.e., a mixture of CH₄ and CO₂ (Ungerfeld, 2014). The synthesis of acetate and butyrate involves the formation of reduced co-factors that require re-oxidization by methanogens (Ungerfeld, 2015). Conversely, propionate fermentation, which is the main precursor of glucose biosynthesis in ruminants, competes with methanogenesis for H₂ (Janssen, 2010). The release of CH₄ is an energetic loss for the animal and a major factor in global climate change (Martin et al., 2010; Opio et al., 2013).

Our knowledge of rumen microbiology has been accumulated by employing culture-based techniques. Studies on isolated cultures have provided a basic understanding of the biochemistry of the isolated strains (Bryant and Burkey, 1953; Hungate, 1960, 1966; Hobson, 1969). Unfortunately, most of the microbial strains in complex habitats cannot be cultivated in pure cultures; therefore, these approaches are of little help when the goal is the elucidation of the relationships between community members. The advent of nucleic acid-based molecular technologies opened up a new culture-independent perspective in microbial ecology. The majority of microbial identification studies rely on 16S ribosomal RNA (rRNA) gene sequencing. The 16S rRNA gene exists across bacterial and archaeal taxa and contains both highly variable and conserved regions (Woese and Fox, 1977; Skillman et al., 2006; Case et al., 2007; Wu et al., 2012),

making it suitable for phylogenetic analyses (Case et al., 2007; Rajendhran and Gunasekaran, 2011). Nevertheless, 16S rDNA sequencing has some drawbacks as it suffers from biased choice of primers, PCR errors, and does not give functional information (Oulas et al., 2015). The meta-omic approaches give a much broader genomic and functional profile generated directly from environmental samples (Venter et al., 2001, 2004; Tyson et al., 2004). “Next generation sequencing” (NGS) has emerged, which employs various chemical reactions for the rapid determination of DNA sequences (MacLean et al., 2009; Metzker, 2010; Goodwin et al., 2016), produces huge datasets from relatively short sequence fragments and uses sophisticated bioinformatics to analyze the results (Huson et al., 2007, 2011; Raes et al., 2007). Initially, the most widely used NGS method in the identification of rumen communities and its functional mechanisms was based on 454-pyrosequencing (McCann et al., 2014; Weimer, 2015; Güllert et al., 2016; Kala et al., 2017). Nowadays, other NGS approaches like IlluminaTM, and Ion TorrentTM have also been employed to investigate these communities (Singh et al., 2014; Mao et al., 2015; Pitta et al., 2016; Comtet-Marre et al., 2017).

At low resolution, the ruminal community appears quite stable (Russell and Rychlik, 2008). Studies have pointed out that diet is the major factor, which has more of an influence than animal species on the rumen microbiome (Jami and Mizrahi, 2012; Henderson et al., 2015; Kala et al., 2017). Nevertheless, there is still limited information available about the core rumen microbiome, i.e., microbes that are shared between individual animals, and about the common functional elements comprising the metabolic networks (Shi et al., 2014; Henderson et al., 2015; Li and Guan, 2017). The aim of this study is to contribute to our understanding of these aspects by examining 10 individual Holstein dairy cow whole metagenome and transcriptome datasets.

MATERIALS AND METHODS

Animals

Ten multiparous cows in the second to fourth lactation period were sampled. The dairy cows were of the Holstein breed, selected from the herd maintained at the Pilot Farm, Faculty of Agriculture, University of Szeged, Hódmezővásárhely, Hungary. This study was carried out in accordance with the recommendations of Hungarian Law of Animal Protection and Charity XXVIII/1998 and Government Decree 40/2013 (II.14.) about animal protection. The protocol was approved by the Ethical Committee of the Faculty of Agriculture, University of Szeged. The best practice veterinary care has been followed and informed consent has been granted by the University of Szeged. The animals were fed with a mixed forage-concentrate diet, using a forage wagon, and water *ad libitum*. **Supplementary Table S1** shows the composition of the diet.

The stomach tube method was used to obtain rumen samples. The plastic collecting tube was 25 mm in diameter and was operated by manual aspiration of rumen fluid. The first 300 mL was discarded to avoid possible contamination with saliva. Sterile

plastic containers were filled with 300 mL of rumen contents, flushed briefly with CO₂ gas, snap-frozen in liquid nitrogen and transported to the laboratory within 1 h. The samples were stored at -20°C. After thawing, the material was filtered through a sterile mesh to remove suspended solids and used for further processing.

DNA and RNA Isolation for Meta-Omic Studies

For total community DNA isolation, 2 mL of rumen liquid samples were used. DNA extractions were carried out using a slightly modified version of the Zymo Research Fecal DNA kit (D6010, Zymo Research, Irvine, CA, United States). The lysis mixture contained 100 µL CTAB (cetyltrimethylammonium bromide) to improve the efficiency (Wirth et al., 2015a,b). After lysis (bead beating), the Zymo Research kit protocol was followed.

For total RNA isolation, 1 mL of rumen liquid samples were taken. The RNA extractions were carried out with the Zymo Research Soil/Fecal RNA kit (R2040, Zymo Research, Irvine, CA, United States). After lysis (bead beating), the Zymo Research kit protocol was followed. The DNA contamination was removed by Thermo Scientific Rapidout™ DNA removal kit (K2981, Thermo Fisher Scientific, Waltham, MA, United States).

The quantity of DNA and RNA was determined in a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, United States) and a Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, United States). DNA purity was tested by agarose gel electrophoresis and on an Agilent 2200 TapeStation instrument (Agilent Technologies, Santa Clara, CA, United States). The quality of the RNA preparation was checked by agarose gel electrophoresis (data not shown).

Next Generation Sequencing and Data Handling

In sample preparation for total metagenome and metatranscriptome sequencing, the recommendations of the Ion Torrent PGM™ sequencing platform were followed (Life Technologies, United States) using Ion Torrent PGM™ 316 chips. Before metatranscriptome sequencing, rRNA was depleted from metaRNA by using the Gram+/Gram- depletion kit in 60:40 ratio (RiboMinus A15020 Life Technologies, United States).

The sequences which were produced by Ion Torrent PGM were denoised, normalized and sequencing artifacts were removed by MG-RAST software pack (DRISSE, Dinamic Trim) (Cox et al., 2010; Meyer et al., 2011; Keegan et al., 2012). The filtered data were downloaded and were further analyzed by Diamond applying default LCA (Lowest Common Ancestor) algorithm (Buchfink et al., 2014). Diamond was set as follows: Blast Mode: BlastX, Min Score: 50, Max Expected: 0,01, Top Percent: 10, Min Support: 1. The data were further analyzed by Megan6 (Huson et al., 2007, 2011, 2016). The NCBI Taxonomy database was used for taxonomic alignment, which is a standard nomenclature and classification repository for the International Sequence Database Collaboration (INSDC) comprising the

GenBank, ENA (EMBL) and DDBJ databases (Federhen, 2015). For functional annotations, the InterPro database was applied. This database integrates several pieces of information about protein families, domains and functional sites (Finn et al., 2017). For carbohydrate active enzyme identification, the CAZy database was used¹ (Lombard et al., 2014). Computing core microbiome and functions Megan6 co-occurrence plot function was applied with the following parameter sets: Threshold (%): 0.01, Min Prevalence (%): 0 Max Prevalence (%): 100, and Probability (%): 100. For functional statistic calculations, the R software was used². The metatranscriptomic sequences were used to link functions to taxa by exporting the data from Megan and further analysis by the R software. The detailed sequence fragment parameters are summarized in **Supplementary Table S2**. The raw data were deposited in NCBI SRA database under the submission SRP148947.

RESULTS

Metagenomic Profiles

Sequencing genomic rumen DNA by Ion Torrent PGM produced an average of 305,884 reads with an average length of 168 bp. In average, 286,786 reads passed the quality control performed by the MG-RAST software package. These were subjected to quality control and were further analyzed by Diamond and Megan6. An average of 106,809 reads was obtained as known taxonomy units based on NCBI Taxonomy. Taking into account that 10 rumen samples were involved in the analysis, the overall read number exceeded 1 million (**Supplementary Table S2**). The most widespread phyla among *Prokaryota* are *Bacteroidetes*, *Firmicutes*, *Proteobacteria*, *Euryarchaeota*, and among *Eukaryota* are *Ciliophora*, *Ascomycota*, and *Neocallimastigomycota*. The taxonomic distribution at class level showed that the most abundant classes within *Bacteria* were *Bacteroidia* followed by *Clostridia*, *Bacilli*, *Spirochaetia* and *Gamma proteobacteria* (**Figure 1**). Within *Archaea*, the class *Methanobacteria* prevails. *Litostomatea*, *Saccharomycetes* and *Neocallimastigomycetes* are the most abundant classes among *Protozoa* and *Fungi*. Rarefaction analysis was performed at genus level. The curves reach its asymptotic nature, indicating that the sequencing depth at this scale is adequate (**Supplementary Figure S1**).

Core Microbiome of the Rumen

Hundreds of microbial genera were identified after the double filtering approach, corroborating that the rumen microbiome is indeed a very diverse community. Co-occurrence analysis was used both on DNA and RNA data to select the planktonic core microbiome, i.e., the group of microbes found in all 10 rumen fluid samples, which were metabolically active. The core included 48 genera (**Figure 2**), which comprised about 18 and 23% of all genera, based on DNA and RNA sequences (**Supplementary Figure S2**).

The class *Bacteroidia* of the phylum *Bacteroidetes* contributed the highest number of core DNA sequences, although the

¹<http://www.cazy.org>

²<https://www.r-project.org>




FIGURE 1 | Normalized distribution of metagenome reads among *Bacteria* classes. Although the absolute number of reads (**Supplementary Table S2**) varied among individual rumen samples (R1–R10), the normalized values indicate a fairly consistent distribution of major taxa.

species diversity of this group was relatively low. *Prevotella* prevailed over other *Bacteroidia* genera. The genus *Bacteroides* was among the top ten abundant genera in this class. The phylum *Bacteroidetes* contained unknown and uncultured species, like *bacterium* P3, P201, F082, F083 or the uncultured class *Prevotellaceae* (Stevenson and Weimer, 2007) (**Figure 2**). *Alistipes*, *Parabacteroides*, *Paraprevotella*, and *Porphyromonas* were less frequent in the class *Bacteroidia* based on DNA data. Nevertheless, it should be noted that from these genera *Parabacteroides*, *Paraprevotella*, and from the uncultured and unclassified species *bacterium* P3, P201 and F082 showed high metabolic activity in RNA data.

The class *Clostridia* (phylum *Firmicutes*) produced fewer sequences in the core but was the most diverse higher taxon in the rumen fluid. Considerable amount of DNA sequence reads pointed at representatives of the genera *Ruminococcus*, *Eubacterium*, *Clostridium*, *Butyrivibrio*, *Oribacterium*, and *Sarcina*. Other genera such as *Blautia*, *Lachnoclostridium*, *Faecalibacterium*, *Roseburia*, *Mogibacterium*, *Ruminoclostridium*, *Oscillibacter*, and *Dorea*, were detected at a lower abundance. The representatives of genera *Butyrivibrio*, *Oribacterium*, and *Lachnoclostridium* had lower apparent metabolic activity than their DNA data suggested. Unclassified sequences were also found, which belonged in the families *Ruminococcaceae*, *Lachnospiraceae* and the phylum *Clostridiales*. It is noteworthy that more than 13 genera of the core 48 taxa belonged in the class *Clostridia* (**Figure 2**).

Also belonging in the phylum *Firmicutes*, the class *Erysipelotrichi* contributed to the planktonic core rumen microbiome with unclassified *Erysipelotrichaceae*, although this class was represented in relative small numbers. The genera *Lactobacillus*, *Streptococcus*, *Bacillus*, and *Paenibacillus*, in the class *Bacilli*, and the genus *Mycoplasma* (class *Mollicutes*) were barely detectable in the DNA based core microbiome. Few sequences of the genera *Succiniclaticum* and *Selenomonas*, cataloged in class *Negativicutes*, from which *Selenomonas* had a few related functions in the functional core compared to the DNA based hits (see Metatranscriptomic Profiles).

The class/phylum *Spirochaetes* was represented by the highly abundant genus *Treponema*.

The genus *Fibrobacter*, belonging in the phylum *Fibrobacteres*, class *Fibrobacteria*, and genus *Bifidobacterium* in the phylum/class *Actinobacteria* were also identified within the core genera. Surprisingly, significantly less RNA sequences mapped to the genus *Fibrobacter* than in the DNA data. Among the minor components of the functionally active core microbiome were other members of *Actinobacteria*, such as the genera *Olsenella*, and *Slackia*.

From the phylum *Proteobacteria*, class *Gammaproteobacteria*, the genera *Acinetobacter* and *Ruminobacter* were identified. The genus *Chlamydia* (class/phylum *Chlamydiae*) was observed in diminishing quantity in DNA reads, and it had even smaller number of RNA hits.

About 1% of all the identified sequences belonged in the phylum *Euryarchaeota*, class *Methanobacteria*. Only the genus *Methanobrevibacter* was verified from this taxon among the core microbes. It is noteworthy that, a comparable number of RNA and DNA sequences were mapped to the genus *Methanobrevibacter*.

The protozoa content of the rumen was about two orders of magnitude lower in cell number than the bacterial cell number (~0.002%). The genus *Entodinium* (protozoan phylum *Ciliophora*, class *Litostomatea*) found its place in the DNA based planktonic core microbiome.

Metatranscriptomic Profiles Metabolic Pathways

Ion Torrent PGM RNAseq produced an average of 680,886 reads with an average length of 185 bp. In average, 661,310 reads passed the MG-RAST software quality control (**Supplementary Table S2**). From the mRNA sequences, in average 296,618 contained predicted protein coding regions with known functions, based on the InterPro database.

Most of the reads were related to metabolic or electron-proton exchange activity. Enzymes involved in biosynthetic, small molecule metabolism, carbohydrate metabolism and




FIGURE 2 | The abundance distribution of the 48 genera comprising the core microbiota calculated from the average DNA-based metagenome and RNA-based metatranscriptome datasets. Error bars indicate the variances among the individual 10 rumen samples.

redox activities were among the most frequent. Housekeeping functions like translation, transport, RNA, DNA and amino acid metabolism were less persistent. The multifaceted picture

was confirmed when putative protein sequences derived from the mRNA data set were projected onto KEGG metabolic pathways providing an integrated picture of global cell




FIGURE 3 | Glucoside hydrolase enzyme families identified in the DNA and RNA datasets, respectively. The inset indicates the proportion of GH sequences relative to the total read numbers.

metabolism. The metabolic pathways showed that in the rumen fluid planktonic microbiome, the metabolism of carbohydrates and volatile fatty acid biosynthesis predominated (**Supplementary Figures S3–S6**).

CAZymes

A great number of cellulases, hemicellulases, and oligosaccharide-degrading enzymes were identified when the filtered reads were probed against the CAZy database. The share of putative CAZymes was 4% in the total RNA database (**Figure 3**). It is noteworthy that DNA-based metagenome data gave similar results, with 6% of DNA reads coding for carbohydrate-degrading genes (**Figure 3**). Moreover, the order of identified enzyme families showed similar tendency. Differences habitually appeared in the relative numbers of corresponding genes; the DNA sequences gave more matches compared to the RNA data. This may be explained by the difference in the number of coded and expressed genes. It is conceivable that not all of the genes in the genome are expressed at all times. Nevertheless, similar trends in the two sets of data clearly indicate that metagenomic DNA data, obtained by NGS, can provide useful information on the main metabolic functions. Among the CAZymes, five glycoside hydrolase (GH) families were prevalent among cellulases in our rumen samples, i.e., GH3, GH5, GH8, GH9, GH44, and GH48. The GH3, GH5, and GH9 families were the most copious among cellulases (**Figure 3**). The GH3, GH5 and GH8 families are widely distributed in bacteria and fungi and have a variety of hydrolyzing activities. The GH9 and GH44 families mainly contain endoglucanases, while GH48 is a reducing end-acting cellobiohydrolase. Eight GH families (GH3, GH8, GH10, GH11, GH16, GH28, GH30, GH43, GH51, GH53, GH67, and GH78) are primarily hemicellulose degraders, and have both endo-acting and debranching enzymes. The identified frequent oligosaccharide-degrading GH families are as follows: GH1, GH2, GH13, GH27, GH29, GH31, GH35, GH36, GH38, GH42, and GH77. GH1, GH2, GH13, and GH77. Carbohydrate-binding modules (CBMs) are specialized

domains in plant cell wall polysaccharide-degrading enzymes. They promote the intimate attachment of catalytic domains to the substrates and enhance the activity of the enzymes³. Nine CBM families, CBM2, CBM3, CBM6, CBM13, CBM20, CBM26, CBM32, CBM34, and CBM48, were identified in both the DNA and mRNA databases. CBM2, CBM3, and CBM6 are specific for cellulose, CBM20, CBM26, and CBM34 have a preference for starch and CBM13, CBM32, and CBM48 families have oligo- and monosaccharide-binding functions. **Supplementary Table S3** contains the entire list of the identified CAZymes.

Core Metabolic Functions

Functional analysis of whole transcriptome data pinpointed thousands of active gene transcripts, confirming the metagenomic data from a distinct view, i.e., the rumen microbiome is a highly active metabolic system that is cumulatively equipped with copious numbers of biochemical functions. Co-occurrence analysis was performed to filter out the most common functions present in our samples. Filtering of the mRNA sequence database yielded at least 587 genes, which may form the core functions indicated by their interactions and presence in all data sets (**Supplementary Figure S7** and **Supplementary Table S4**). According to the InterPro database, 62 genes from this pool apparently code for enzymes involved in metabolic pathways (**Figure 4**). This is a small number for the enzymes to comply with the requirements of basic biochemical life-sustaining fermentative pathways, which is likely due to the stringent filtering conditions used in the co-occurrence analysis. The remaining transcripts are housekeeping genes required for the maintenance of basic cellular functions. The core metabolic processes provided satisfactory coverage of the entire microbiological food chain, from carbohydrate metabolism to methanogenesis. From the 62 core metabolic function coding genes, 18 belonged to carbohydrate processing, which comprised

³<https://www.cazypedia.org>




FIGURE 4 | The distribution of the core microbial functions calculated from the average RNA-based metatranscriptome dataset. Error bars indicate the variances among the individual 10 rumen samples. The inset indicates the relative partition of the functional groups within the metatranscriptome.

26% of all core transcripts. Most of them encoded glycoside hydrolase families.

Among small molecule metabolic processes, pyruvate phosphate dikinase (PPDK) prevailed, which is a central glycolytic enzyme, together with phosphoenolpyruvate (PEP) synthase (PPsTK) (Berman and Chon, 1970; Imanaka et al., 2006). The second most frequently identified sequence indicated ATP-utilizing phosphoenolpyruvate carboxykinase (PEPCK), which participates in gluconeogenesis in co-operation with malate L-lactate dehydrogenase (Madern, 2002). Uronate

isomerase is an enzyme involved in the intramolecular rearrangement of glucose (Williams et al., 2006). Folypolyglutamate synthase might be implicated in the biosynthesis of the important methanogen intermediary tetrahydrofolate (Bognar et al., 1985). Additional enzymes found in the core functions complementing the tetrahydrofolate pathway include tetrahydrofolate dehydrogenase/cyclohydrolase (Pomper et al., 1999) and formate-tetrahydrofolate ligase (FTHFS) (Chistoserdova et al., 1998) (Figure 4 and Supplementary Figure S4).

A diverse group of redox enzymes such as pyruvate-flavodoxin oxidoreductase (PFOR), glyceraldehyde/erythrose phosphate dehydrogenase family, the alpha subunit of indolepyruvate ferredoxin oxidoreductase, malic oxidoreductase, and L-lactate dehydrogenase indicated highly vigorous redox processes, which made up 28% of all core transcripts. Glutamate dehydrogenase, glutamate synthase and glutamate-5-semialdehyde dehydrogenase within this group signaled active ammonia assimilation (Yuan et al., 2009) by the rumen community. HydE, responsible for the maturation of [FeFe], hydrogenase suggests the presence of hydrogen metabolism in each rumen sample (**Figure 4** and **Supplementary Figure S6**).

A distinct group of the represented core functions take part in the synthesis of precursor metabolites and energy supply including ATP dependent 6-phosphofructokinase, fructose-1,6-bisphosphate aldolase, phosphoglycerate kinase, enolase, phosphoglucose isomerase, adenylate kinase and pyruvate kinase were found in high transcript numbers. Numerous additional enzymes were identified in the core functional group, which are also implicated in various biosynthetic pathways (**Figure 4**).

Overall, 7% of the total transcripts represented enzymes playing a part in organic acid metabolic processes, i.e., acetate/propionate kinase, methylmalonyl-CoA, acetyl-CoA carboxylase carboxyl transferase, succinate CoA transferase and butyrate kinase (**Figure 4** and **Supplementary Figure S4**).

Although they represent only 1% of the total core transcripts, Archaea enzymes performing methanogenesis play important functions in maintaining the proper redox balance in the rumen. Methyl-coenzyme M reductase alpha, beta, and gamma subunits and 5,10-methylenetetrahydromethanopterin reductase were detected in all of the rumen samples investigated (**Figure 4** and **Supplementary Figure S6**).

Linking Core Functions to Taxa

In most metagenomic studies, the composition of the microbial community is determined and changes in relative abundances are recorded as the results of changes in various environmental conditions or feeding regimes. More recently, important experiments have complemented this information aiming at functional aspects by metatranscriptome, proteome and metabolome investigations concentrating on the “who does what?” question. The metabolic functions are grouped according to the InterPro classification⁴.

In all of the functional groups associated with *Bacteria*, the genus *Prevotella* (phylum *Bacteroidetes*) prevailed (**Figure 5**), due to its vast predominance in the rumen planktonic microbiota and the versatile metabolic pathways possessed by this genus. Accordingly, the share of *Prevotella* in carbohydrate (~63%), small molecule (~50%), organic acid (~53%) metabolisms, oxidation-reduction processes (~44%), and generation of precursor metabolites (~42%) was high, indicating that the majority of the metabolic “work” taking place in the rumen fluid is carried out by representatives of this taxon (**Figure 5**). Seven additional genera complemented the picture participating in the various metabolic pathways with varying activity. These

were as follows: *Bacteroides*, *Ruminococcus*, *Bifidobacterium*, *Clostridium*, *Lactobacillus*, *Paraprevotella* and *Eubacterium*. It is interesting to note that the diversity ranking at the genus level is different from the abundance distribution based on the metagenomic data (**Figure 2**). The divergences between DNA and RNA based taxonomy data suggest that abundance and functional importance are not necessarily coupled.

The genus *Bacteroides* extensively contributed to carbohydrate, small molecule and organic acid metabolisms and played a significant role in other Bacteria-linked metabolic processes as well. The genus *Ruminococcus* was primarily active in the generation of precursor metabolites and energy although these species also participated in other bacterial tasks. *Clostridium* was more active in handling oxidation-reduction processes than *Ruminococcus*, and lagged behind *Ruminococcus* in carbohydrate metabolic process performance. In spite of their low relative abundance in the community, the genus *Lactobacillus* displayed remarkable contribution to tasks related to the generation of precursor metabolites and energy. *Bifidobacterium* showed considerable involvement in carbohydrate metabolism-related pathways and a noticeable one in organic acid metabolism.

Paraprevotella and *Eubacterium* were detected in the carbohydrate and small molecule metabolic processes. In the functional group “oxidation- reduction processes,” an uncultured bacterium was observed as the second dominant genus.

The genera *Kandleria*, *Parabacteroides*, *Porphyromonas*, *Selenomonas*, *Alistipes*, *Catenibacterium*, *Barnesiella*, and *Mitsuokella* were found less frequently among the various metabolic functions, perhaps due to their low abundance in the planktonic rumen microbiota. *Lachnospiraceae* and *Treponema* were the apparent exceptions as they were present at a relatively high number in the core rumen microbiome and proportionally low related activity.

In the methanogenesis-related core functions, the picture is completely different from the other functional groups, involving only archaeal taxa. In this function, the genus *Methanobrevibacter* predominated (~88%). An uncultured methanogenic archaeon was also detected as the second most active in methanogenesis (~6%). The contributions of the genera *Methanobacterium* (~2%), an incompletely described archaeon genus *Candidatus* (~2%), *Methanomassiliicoccales* (~2%), *Methanosphaera* (~1%), *Methanosarcina* (~0.5%), and *Thermoplasmatales* (~0.5%) were negligible.

DISCUSSION

The Core Microbiome

The following discussion is restricted to genera present in the core microbiome and have functions associated to them (**Supplementary Table S4**). This information will be related to the results of previous rumen studies. In this study, our fundamental assumption has been that the taxa occurring together in the various rumen microbiota have shared functional connections and are therefore in some sort of metabolic relationship. This may not be the case in every individual interacting partner pair, but it is a reasonable postulation to find

⁴<https://www.ebi.ac.uk/interpro/>




FIGURE 5 | Coupling functional groups to taxonomic genera determined from the metatranscriptomic dataset.

interrelationships between taxonomic and functional analyses in the core community.

Within the domain *Bacteria*, the majority of the identified sequences were annotated to the genus *Prevotella* (class *Bacteroidia*) (Figure 2 and Supplementary Figure S2). Previous studies, using qPCR and/or 16S rDNA sequencing, also identified this genus as the most abundant in the rumen bacterial population (Stevenson and Weimer, 2007; Jami and Mizrahi, 2012). Further confirmation came from the analysis of the Sus-like (Starch utilization system) polysaccharide utilization loci (Naas et al., 2014; Rosewarne et al., 2014; Güllert et al., 2016). Changing dietary conditions did not alter *Prevotella* abundance (Stevenson and Weimer, 2007; Bekele et al., 2010; Purushe et al., 2010; Pitta et al., 2014; Lyons et al., 2017), although an age-related shift from *Prevotella* to *Succinivibrio* was noted recently (Liu et al., 2017). This is likely due to their ability to serve the wellbeing of their host in various ways, exploiting their high degree of genetic diversity (Avgustin et al., 1994; Ramšak et al., 2000) and remarkable metabolic versatility (Bryant et al., 1957;

Wen et al., 1996; Matsui et al., 2000). In addition, *Prevotella* is a highly diverse taxon comprising various functional niches in different systems (Purushe et al., 2010).

The second most widespread group in the class *Bacteroidia* was the genus *Bacteroides*. This genus, like *Prevotella*, can utilize polysaccharides as an energy source. Apparently, some rumen *Bacteroides* can also decompose cellulose (Naas et al., 2014). Smaller genera in the class *Bacteroidia* include *Parabacteroides*, *Paraprevotella* and *Porphyrmonas*, in addition to the *uncultured Prevotellaceae* family. These are usually detected in cow manure and their relative abundances change during the transition from developing and mature rumen (Dowd et al., 2008; Wu et al., 2012).

The classes *Bacteroidia* (phylum *Bacteroidetes*) and *Clostridia* (phylum *Firmicutes*) together represented ~47–30% of the rumen core microbiome. In a recent study, the ratio of the two taxa in the whole rumen community was close to 1:1 (Güllert et al., 2016). Interestingly, in the biogas producing anaerobic microbial community metabolizing similar

lignocellulosic substrates, *Clostridia* surpass greatly *Bacteroidia* (Güllert et al., 2016; Bozan et al., 2017). Apparently, the two classes differ in their colonization strategies. Only a few genera of the class *Bacteroidia* were present in the rumen, although in large relative abundances, whereas a remarkable diversity of genera characterized the class *Clostridia*.

Among *Clostridia*, the thoroughly studied genus *Ruminococcus* was frequently found. Their interactions with other species may make this genus more important than their abundance implies (Fondevila and Dehority, 1996; McSweeney et al., 1999; Koike and Kobayashi, 2009; Christopherson et al., 2014; Dai et al., 2015). Members of the family *unclassified Lachnospiraceae* and the genera *Eubacterium* and *Blautia* are among the taxa in the core of the cow rumen and present also in the kangaroo forestomach (Henderson et al., 2010; Godwin et al., 2014). These bacteria acquired the Wood-Ljungdahl pathway, which is alternatively called the reductive acetyl-CoA or the reductive acetogenesis pathway (Hattori, 2008; Ragsdale and Pierce, 2009; Xu et al., 2009; Gagen et al., 2015; Kelly et al., 2016) (**Supplementary Figure S4**).

The genus *Clostridium* possess high cellulolytic activity and actively produces H_2 (Lin et al., 2007). The partial pressure of H_2 determines the rate of methanogenesis and the assortment of short chain fatty acids generated in the rumen (Hegarty and Gerdes, 1999; Janssen, 2010). Methanogens outcompete the microbes involved in reductive acetogenesis because acetogens are less efficient at obtaining energy from the oxidation of H_2 (Le Van et al., 1998; Siri Wongrungson et al., 2007; Ungerfeld, 2015). The genus *Butyrivibrio* is present in a large selection of ruminants and has been characterized as an oligosaccharide-degrading bacterial taxon (Dai et al., 2015; Henderson et al., 2015).

The genus *Oribacterium* was observed as a particle associated bacterium in the rumen and the genus *Mogibacterium* was identified as a predominant member of the cattle gastrointestinal tract, although little is known about their functions (Mao et al., 2015; Schären et al., 2017). The genera *Faecalibacterium*, *Roseburia*, *Ruminiclostridium*, *Oscillibacter*, and *Dorea* were represented in low abundance. These diverse genera are active in cellulose and hemicellulose decomposition and ferment various sugars to short chain fatty acids (Duncan et al., 2006; Flint et al., 2008; Koike and Kobayashi, 2009; Thoetkiattikul et al., 2013; Ravachol et al., 2015).

The genera *Sharpea*, *Lactobacillus*, *Streptococcus*, *Bacillus*, and *Paenibacillus* represented the core members of the class *Bacilli* (phylum *Firmicutes*). Perhaps not surprisingly, the lactic acid bacteria *Streptococcus* and *Lactobacillus* tolerated aciduric challenge well in the cow rumen (Petri et al., 2013). Representatives of the genera *Bacillus* and *Lactobacillus* produce organic acids, some of which may endow probiotic effects (Qiao et al., 2010; Seo et al., 2013; Ushakova et al., 2013). The genus *Paenibacillus* has been noted for its hemicellulose-degrading activity (Kala et al., 2017), while the genus *Sharpea* tends to form and metabolize lactate and thereby increased its relative abundance in low methane yield sheep rumen (Kamke et al., 2016).

Within the core rumen planktonic microbiome, the genera *Succiniclasicum* and *Selenomonas*, were also present, belonging

to the class *Negativicutes* (phylum *Firmicutes*), although in low abundance. The genus *Selenomonas* is lactate-utilizing bacteria (Hackmann and Firkins, 2015). *Succiniclasicum* members apparently differed in quantity between hay diet- and high grain diet-fed goats and have been described as starch degraders (Kim et al., 2011; Huo et al., 2014).

Three identified taxons in the core microbiome belonged in the class *Actinobacteria* (phylum *Actinobacteria*), these were the genera *Bifidobacterium*, *Olsensella*, and *Slackia*. Certain members of these genera are considered probiotic in both ruminants and humans as they metabolize oligosaccharides and release lactic acid, which helps to control the normal microflora (Picard et al., 2005; Nagai et al., 2010; Kamke et al., 2016).

The genus *Treponema* belong to the class/phyla *Spirochaeta*. Microbes in this genus are important fatty acid producers in the rumen (Yamada and Yukphan, 2008; Henderson et al., 2010; Li et al., 2012). The genus *Fibrobacter*, member of the class *Fibrobacteria* (phylum *Fibrobacteres*), contains well-known cellulose degraders (Dai et al., 2015). Our transcriptomic data indicate that the activity of this genus is lower than suggested by the DNA based hits. Similar phenomenon could be observed in case of the genera *Sharpea*, *Treponema*, *Butyrivibrio*, *bacterium P3*, *bacterium P201* and *Lachnoclostridium*. This may imply that, not always the predominant genera play essential role and some microbes present in small numbers may have significant activity in the microbial ecosystems.

In the rumen core microbiome the single genus *Methanobrevibacter* was found as the most prominent *Archaea* taxon. They are hydrogenotrophic methanogens and reduce CO_2 to CH_4 when H_2 is present as an electron donor (Deppenmeier et al., 1996). The high rate of short chain fatty acid passage through the rumen wall via active uptake may explain the predominance of hydrogenotrophic *Archaea*, as the short chain fatty acid (SCFA) uptake is faster than the growth rate of acetate-utilizing methanogens (Dijkstra, 1994).

The total *Protozoa* and *Fungi* cell number in the rumen is two orders of magnitude lower than that of the *Bacteria*. In our data among *Protozoa*, the single genus *Entodinium* (class *Litostomateae*) positioned itself in the DNA based core microbiome, which reflects the uneven distribution of *Protozoa* in the individual rumen microbiota. The genus *Entodinium* is very efficient in starch fermentation and is a frequently detected protozoan in the rumen. (Jouany and Ushida, 1999; Zhang et al., 2017). Surprisingly, none of the *Fungi* were present in the DNA and RNA based core microbiome, suggesting that the fungal community is extremely varied across the individual rumen communities.

Core Functions and Their Associated Microbes

Numerous studies recognized that the functional and phylogenetic distribution of microbes in the rumen comprise an integrated system; therefore, both aspects should be investigated together to better understand the complex metabolic processes taking place in this ecosystem (Deusch et al., 2015; Bielak et al., 2016; Mayorga et al., 2016; Shabat et al., 2016; Wang et al., 2016; Shen et al., 2017). In the present study, core metabolic functions

and their accompanying microbes were determined from double filtered metatranscriptomic data by pairing the functional groups with the most probable taxonomic units. This resulted in a comprehensive coverage of the microbial food chains from complex carbohydrates to end products, i.e., SCFA and CH₄, to be utilized or released by the host animal. In the following section, we therefore attempt to reconstruct a metabolic map of the physiological events taking place in the cattle rumen (**Figure 6**) and discuss the identified microbes associated with these biochemical reactions, extending the findings of similar previous studies (Stevens and Hume, 1998; Russell and Rychlik, 2008; Ungerfeld, 2014, 2015; Jiang et al., 2016; Deusch et al., 2017).


A great diversity of carbohydrate-active enzymes (CAZy) take part in the breakdown of the lignocellulosic feed components (Ferrer et al., 2012; Pope et al., 2012; Dai et al., 2015; Comtet-Marre et al., 2017; Kala et al., 2017; Svartström et al., 2017; Stewart et al., 2018). Our data corroborated these observations and numerous hemicellulose-, cellulose- and oligosaccharide-degrading enzymes, collectively called glycosyl hydrolases (GH), were detected in the first steps of the core energy providing metabolic pathways (**Figure 6**). GH8, GH28, GH30 and GH43 represent the hemicellulose degrading enzymes with polygalacturonase, xylanase and xylosidase activities. The genera *Prevotella* and *Bacteroides* predominate among microbes expressing these GH families (**Supplementary Table S3**). In the case of GH30, the genus *Prevotella* prevailed (~30%), but other members of the class *Bacteroidia* also exhibited significant GH30 activity. The core cellulases might be GH8 and GH9. It is noteworthy that the GH9 family has been recognized before in numerous rumen ecosystems, indicating an important and general role in cellulose degradation (Brulc et al., 2009; Hess, 2011; Wang et al., 2013; Dai et al., 2015; Shinkai et al., 2016; Comtet-Marre et al., 2017). GH9 could be assigned primarily to the genus *Prevotella* (~30%), although the genera *Bacteroides* (~16%), *Ruminococcus* (~12%), *Clostridium* (~2%) and *Fibrobacter* (~0.1%) expressed transcripts belonging in this enzyme family. Apparently, the oligosaccharide-degrading enzymes had the most diverse range of catalysts for carbohydrate breakdown. GH77 has amylomaltase and 4- α -glucanotransferase (EC 2.4.1.25) activity, which are mainly provided by the genera *Prevotella* (~65%), *Ruminococcus* (~3%) and *Clostridium* (~3%). This GH family was found in the buffalo rumen among the highly active oligo-GH families (Patel et al., 2014). *Prevotella* and *Bacteroides* were the predominant genera harboring the enzyme families GH31 (*Prevotella*: ~70%, *Bacteroides*: ~11%), GH36 (*Prevotella*: ~48%, *Bacteroides*: ~8%), and GH27 (*Prevotella*: ~50%, *Bacteroides*: ~14%). Alpha-glucuronidases play an essential role in the complete hydrolysis of hemicellulose and were studied in *Bacteroides vulgaris* and *Dysgonomonas mossii* (Lee et al., 2012). The enzyme family GH67 was linked to the genera *Chryseobacterium* (~29%), *Dysgonomonas* (~10%) and *Paludibacter* (~8%).

Taking the transcriptomic and taxonomic data together, the genera *Prevotella*, *Bacteroides* and *Ruminococcus* seem to predominate in the deconstruction of the lignocellulose rich feed.

Previous reports highlighted that, in the rumen microbiota, the phyla *Firmicutes* and *Bacteroidetes* are almost equally responsible for cellulose degradation, while in hemicellulose decomposition the members of *Bacteroidetes* predominated (Comtet-Marre et al., 2017). Our data showed that *Prevotella* was primarily responsible for carbohydrate metabolism (**Supplementary Table S3**). *Prevotella* provided fewer transcripts of the cellulose GH9 family (~30%) than those of the GH77 (~65%), GH31 (~70%), GH28 (~57%), and GH43 (~90%) enzymes, which are associated with oligosaccharide-degradation and hemicellulases.

Monomeric hexoses and pentoses are fermented to phosphoenolpyruvate in the subsequent metabolic steps (red boxes in **Figure 6**). Phosphoenolpyruvate is an important intermediary of glycolysis and gluconeogenesis (Parmar et al., 2015). We identified three genes in the core metabolic functions participating in this transformation, i.e., ATP-dependent 6 phosphofructokinase (IPR022953), enolase (IPR000941) and fructose-1,6-bisphosphate class 3 (IPR009164). ATP-dependent 6 phosphofructokinase catalyzed the transfer of a phosphoryl group from ATP; this is an important reaction in a wide variety of biological processes (Hellings and Evans, 1987). This enzyme phosphorylates fructose-6-phosphate to fructose-1,6-bisphosphate; the reaction is a key regulatory step in the glycolytic pathway (Raben et al., 1995; Wegener and Krause, 2002). Our data suggested several taxa as the source of this enzyme activity, including the genera *Prevotella* (~59%), *Bacteroides* (~6%), *Ruminococcus* (~4%) and *Kandleria* (~4%). Enolase (2-phospho-D-glycerate hydrolase) is a homodimer enzyme that catalyzes the reversible dehydration of 2-phospho-D-glycerate to phosphoenolpyruvate in the presence of Mg²⁺ (Lebioda and Stec, 1991). The genera expressing this gene were *Prevotella* (~39%), *Methanobrevibacter* (~6%) and *Lactobacillus* (~5%). Fructose 1,6-bisphosphatase converts fructose 1,6-bisphosphate to fructose 6-phosphate (**Figure 6**). This is a key reaction in gluconeogenesis (Marcus et al., 1986). The genera *Prevotella* (~55%), *Bacteroides* (~14%) and *Eubacterium* (~5%) are the main suppliers of this activity. Phosphoenolpyruvate is converted to pyruvate in glycolysis or to oxaloacetate in the pathway leading to gluconeogenesis. In the reversible reaction, the pyruvate, phosphate dikinase (IPR010121), pyruvate kinase (IPR001697) and phosphoenolpyruvate synthase (IPR006319) participated (Berman and Chon, 1970; Imanaka et al., 2006), and were among the core metabolic functions. The highest number of transcripts found for pyruvate, phosphate dikinase and the related genera are principally *Prevotella* (~68%), *Eubacterium* (~3%) and *Ruminococcus* (~3%). This enzyme catalyzes the reversible conversion of ATP and pyruvate to AMP and phosphoenolpyruvate (Evans and Wood, 1968).

Pyruvate kinase was found in the genera *Lactobacillus* (~15%), *Prevotella* (~15%), *Ruminococcus* (~8%), *Kandleria* (~8%), and *Eubacterium* (~3%). This enzyme plays a role in the final step of glycolysis, i.e., the conversion of phosphoenolpyruvate to pyruvate (Muirhead, 1990). Phosphoenolpyruvate synthase was expressed predominantly by the genera *Prevotella* (~32%), *Phascolarctobacterium* (~26%) and *Acidaminococcus* (~9%). It takes part in gluconeogenesis and converts pyruvate back into



phosphoenolpyruvate (Berman and Chon, 1970). Although the genera *Phascolarctobacterium* and *Acidaminococcus* are important in the pyruvate to phosphoenolpyruvate conversion, they did not fit into the core microbiome because they were not detectable in all rumen samples.

In the rumen, pyruvate is converted anaerobically to SCFA (Stevens and Hume, 1998) (green arrows in **Figure 6**). The first step in this process is the generation of precursor metabolite acetyl-CoA from pyruvate (Charon et al., 1999). According to our transcriptomic data, pyruvate-flavodoxin oxidoreductase (IPR011895) is the key enzyme in this process from *Prevotella* (~51%), *Clostridium* (~13%) and *Bacteroides* (~5%). Another reaction of pyruvate may lead to the simultaneous formation of formate and acetyl-CoA. This enzyme may participate in protecting the bacteria against oxidative stress (Nakayama et al., 2013). Pyruvate formate lyase (PFL) (IPR012838) catalyzes the conversion of pyruvate and CoA to formate and acetyl-CoA has been found in representatives of *Prevotella* (~13%), *Veillonellaceae* (~12%), and *Lachnospiraceae* (~5%). Formate acetyltransferase (IPR005949) (black arrows in **Figure 6**) acts in the reverse direction, and was detected in the genera *Prevotella* (~9%), *Butyrivibrio* (~8%), *Eubacterium* (~6%), *Succinatimonas* (~6%), *Treponema* (~6%) and *Clostridium* (~4%).

Formate in the rumen serves as a precursor metabolite for hydrogenotrophic methanogenesis (black lines in **Figure 6**) or reductive acetogenesis (Ragsdale and Pierce, 2009; Ungerfeld, 2014). In reductive acetogenesis, the formate-tetrahydrofolate ligase (FTHFS) gene was identified among the core metabolic functions (blue line in **Figure 6**). FTHFS (IPR000559) catalyzes the ATP-dependent activation of formate ions via its addition to the N10 position of tetrahydrofolate (Ragsdale and Pierce, 2009). FTHFS is widely considered indicative of the Wood-Ljungdahl

pathway of autotrophic CO₂ fixation (acetogenesis) and was used to identify homoacetogenic bacteria (Xu et al., 2009; Henderson et al., 2010; Müller et al., 2016) (**Supplementary Figure S4**). Interestingly, the gene encoding this enzyme was largely supplied by an uncultured bacterium (~75%), followed by *Bifidobacterium* (~4%), *Bacteroides* (~3%) and *Ruminococcus* (~3%). The 75% “uncultured” suggests that there are still substantial unexplored fields of homoacetogenesis (Campanaro et al., 2016; Müller et al., 2016).

Although the genera *Bifidobacterium* and *Bacteroides* contained the FTHFS gene, the enzyme does not exclusively serve the Wood-Ljungdahl pathway in these microbes; it functions rather as a methyltransferase in purine and glycine degradation and in the metabolism of some sulfate-reducing bacteria (Xu et al., 2009; Henderson et al., 2010). The genera *Ruminococcus*, *Clostridium* (~1%), *Blautia* (~1%) and order *Clostridiales* (~0.5%) include known homoacetogens but they are represented in low transcript frequency (**Supplementary Table S3**).

Along the pathway of heterotrophic acetate production, acetyl-CoA becomes phosphorylated to acetyl-phosphate (Schuchmann and Müller, 2014). The acetate/propionate kinase enzyme (IPR004372) can carry out this reaction. Acetate kinase catalyzes the acetyl-phosphate to acetate reaction with concomitant ATP generation (Hasona et al., 2004). The corresponding transcript was widespread in the genus *Prevotella* (~69%) followed by *Bacteroides* (~7%), *Clostridium* (~3%), *Ruminococcus* (~3%) and *Lactobacillus* (~3%). Acetate kinase can be involved both in the reductive acetogenesis and heterotrophic acetate production (Schuchmann and Müller, 2014); thus, the genera attached to this reaction are probably mixed. Butyrate is produced by butyrate kinase (IPR011245)

from butyryl-CoA (Louis et al., 2004). The corresponding members of the rumen fluid microbial community were *Prevotella* (~52%) and *Bacteroides* (~14%), respectively. Our data indicated that several gene products contributed to the regulation of acetate and butyrate production, i.e., the alpha subunit of acetyl-CoA carboxylase (IPR001095), the beta subunit of acetyl-CoA carboxylase carboxyl transferase (IPR000438) and the acetyl-CoA biotin carboxyl carrier (IPR001249) (green lines in **Figure 6**). These enzymes carry out the carboxylation of acetyl-CoA to malonyl-CoA, thus they tightly regulate the SCFA synthesis (Rock and Jackowski, 2002). The taxa linked to this regulation were the genera *Prevotella* and *Ruminococcus* (**Supplementary Table S3**).

Two gene products were identified in the core functions in the pathway of oxaloacetate biosynthesis. One of them was the phosphoenolpyruvate carboxykinase (IPR001272), which catalyzes the reversible carboxylation of phosphoenolpyruvate to oxaloacetate (purple arrow in **Figure 6**). This has been envisioned as a major carboxylation enzyme for the generation of oxaloacetate intermediate in succinate-producing saccharolytic anaerobes (Samuelov et al., 1991; Schöcke and Weimer, 1997). Our results revealed that the genera *Clostridium* (~23%), *Ruminococcus* (~23%), *Treponema* (~23%) and *Eubacterium* (~6%) contributed to this particular metabolic pathway. An alternative oxaloacetate synthetic pathway from pyruvate also exists. The pyruvate carboxylase (IPR005930) enzyme is essential in this function, shifting the metabolism toward gluconeogenesis (Jitrapakdee et al., 2008). The reaction involves the irreversible ATP-dependent carboxylation of pyruvate to oxaloacetate; we detected the likely presence of this reaction in the genera *Eubacterium* (~17%), *Clostridium* (~17%), *Oscillibacter* (~17%), and *Megasphaera* (~11%), respectively.

Further on in this metabolic route, malate is produced from oxaloacetate by malate L-lactate dehydrogenase (IPR003767). This is more favorable than the reverse reaction in bacterial species (Honka et al., 1990) and is carried out by a diverse community of microbial genera, including *Slackia* (~6%), *Clostridium* (~5%), *Olsensella* (~5%), *Blautia* (~3%), *Escherichia* (~3%), and *Lactobacillus* (~3%). Malate and fumarate are precursor metabolites for succinate (purple arrows in **Figure 6**). Succinate plays very important role in the rumen fermentation (Bryant et al., 1957; Wen et al., 1996; Matsui et al., 2000; Russell and Rychlik, 2008; Purushe et al., 2010; Reichardt et al., 2014; Bozan et al., 2017; Deusch et al., 2017). Our results indicated that the core succinate producers in the rumen were representatives of *Clostridium*, *Eubacterium*, and *Ruminococcus* (Russell and Rychlik, 2008; Reichardt et al., 2014). The ruminal decarboxylation of succinate is rapid and the resulting propionate plays a key role in maintaining glucose homeostasis (Purushe et al., 2010). Propionate is synthesized via the succinyl-CoA, methylmalonyl-CoA, and propionyl-CoA metabolic pathway (purple arrows in **Figure 6** and **Supplementary Figure S5**). The beta chain of methylmalonyl-CoA mutase (IPR004608) (Takahashi-Iñiguez et al., 2012) was identified, indicating that this is part of the core biochemical pathways in the rumen. This enzyme catalyzes the isomerization of succinyl-CoA to methylmalonyl-CoA and the enzyme activity was carried

by the genera *Prevotella* (~68%), *Phocaeicola* (~10%) and *Bacteroides* (~9%). Succinate-CoA transferase (IPR017821) may catalyze the pathway for the decarboxylation of succinate to propionate⁵. The gene coding for this enzyme could be linked to the genera *Prevotella* (~61%), *Phascolarctobacterium* (~13%), and *Bacteroides* (~5%), respectively. The propionate-producing genera *Phocaeicola* and *Phascolarctobacterium* (Al Masalma et al., 2009; Maspolim et al., 2015) did not make it into the core rumen microbiome because some of the rumen samples did not contain these genera in detectable amounts. This implies that the genera *Prevotella* and *Bacteroides* predominate in the metabolic conversion of succinate to propionate.

The principal route to methanogenesis is the hydrogenotrophic one, in which CO₂, formate and H₂ are used as substrates (Hook et al., 2010; Janssen, 2010; Morgavi et al., 2012; Wirth et al., 2012; Ungerfeld, 2015; Wallace et al., 2015, 2017; Kamke et al., 2016) (black arrows in **Figure 6**). Reducing equivalents and CO₂ are released along the fermentation process (Stevens and Hume, 1998; Russell and Rychlik, 2008). Molecular H₂ is produced actively by hydrogenases. We located in the core functions the [FeFe] hydrogenase maturation HydE gene (IPR024021), which is essential for the maturation the [FeFe] hydrogenase (King et al., 2006). The genera *Roseburia* (~16%), *Clostridium* (~14%), *Ruminococcus* (~9%), *Bacteroides* (~8%) and *Megasphaera* (~6%) displayed the corresponding transcripts. Formate, may be used for methanogenesis directly or after splitting it to H₂ and CO₂ by the formate hydrogenlyase complex (Russell and Rychlik, 2008; McDowall et al., 2014; Ungerfeld, 2015). H₂ plays an important role in the overall rumen function and in animal nutrition. Efficient H₂ removal leads to a nutritionally more favorable assortment of SCFA and to an increased rate of fermentation by eliminating the inhibitory effect of H₂ on acetogens (Henderson et al., 2010; Petri et al., 2013; Shi et al., 2014; Mao et al., 2015; Ungerfeld, 2015). Our results further corroborated the dominance of hydrogenotrophic methanogenesis in the rumen. In addition, we identified four essential gene transcripts among the core functions, which were related to methanogenesis, i.e., the alpha, beta and gamma subunits of methyl coenzyme M reductase (IPR016212, IPR003179, IPR003178) and 5,10-methylenetetrahydromethanoprotein reductase (IPR019946). All known methanogens express the enzyme methyl coenzyme M reductase, which catalyzes heterodisulfide formation between methyl-coenzyme M and coenzyme B and the subsequent release of methane (Ferry, 1999; Hallam et al., 2003). In our data set, the hydrogenotrophic genus *Methanobrevibacter* expressed these genes, although we also noted an unknown methanogenic archaeon associated with relatively high expression level (**Supplementary Table S3**). This suggested that the ruminal archaeal community has not been fully explored to date (Campanaro et al., 2016). Also, 5,10-methylenetetrahydromethanopterin reductase, an F420-dependent enzyme of methanogenesis (Brömmelstroet et al., 1990, 1991), was linked in our transcriptomic data to *Methanobrevibacter* (~92%) and *Methanobacterium* (~6%).

⁵<https://www.ebi.ac.uk/interpro/entry/IPR017821>

CONCLUSION

The metagenome and metatranscriptome of the rumen fluid fractions of ten lactating cows showed a fairly consistent picture, which corroborated and extended our knowledge from similar previous studies. Ten separate rumen samples were analyzed, about one third of the average 305,844 Ion Torrent DNA reads could be associated with taxonomic information in the NCBI Taxonomy database. In the RNAseq experiments an average of 680,886 reads were obtained and approximately half of them predicted proteins of known functions. Each set of data was subjected to quality control and filtering before co-occurrence analysis, which was based on the assumption that those microbes which occurred together were likely to form physiological relationships.

A core microbiome, i.e., a collection of taxa present in all ten sequenced samples, was established. The taxonomic core microbiome consisted of 48 genera belonging in *Bacteria* and 1 *Archaeon*. Based on their relative abundances, about a dozen genera formed the majority of the microbiota; the genus *Prevotella* exceedingly predominated the community. *Fungi* were not placed in the core microbiome, which might suggest that either fungi formed a diverse community or they were present at a very low number in our samples and their DNA sequence reads could not reach the threshold.

Thousands of mRNA sequences pointed to active proteins, implicating a complex and diverse metabolism taking place in the rumen. From the 587 functions, a core functional group was distinguished based on co-occurrence analysis, but only 62 of them could be assigned to metabolic functions. This is clearly a low number to cover all possible metabolic pathways and is ascribed to very stringent filtering conditions and to the selection of common elements in 10 separate rumen fluid samples. Nevertheless, even this minimal functional core revealed key enzymes participating in various metabolic processes. As expected, a diverse and rich collection of enzymes was involved in carbohydrate metabolic processes, but other functional groups were also sufficiently represented. Transcripts coding for enzymes active in methanogenesis made up 1% of the core functions.

The genera associated with the core metabolic functions were identified. The main conclusion drawn from this investigation was that for all metabolic functions, performed primarily by *Bacteria*, several genera could provide the necessary activity. In other words, there always seems to be a “back-up” microbial team to substitute the predominant bacteria to accomplish any given metabolic function. Nevertheless, the key actors in most metabolic functions belong to the genus *Prevotella*. The *Prevotella* predominance is partly due to their massive abundance and partly to the metabolic versatility of this taxon. Contrary to the extensive potential to share the contribution to the tasks among bacterial taxa, methanogenesis seemed to have limited possibilities, i.e., only the hydrogenotrophic pathway existed with a limited possible role for “unidentified” methanogens. The potential bypass biochemical routes make the diversion of metabolism toward advantageous pathways by managing the rumen community composition a considerable challenge.

Nonetheless, methane emission mitigation seems conceivable via targeting the hydrogenotrophic genus *Methanobrevibacter*.

AUTHOR CONTRIBUTIONS

KK conceived the study, participated in its design and evaluation. GK and JH collected the rumen samples and took part in the assessment of the data. RW, GM, and BK performed the sequencing experiments and statistically evaluated the metagenomic data sets. KK, RW, ZB, GR, and AS composed the manuscript. All the authors agreed in publishing the final version of this paper.

FUNDING

This study has been supported in part by the Hungarian National Research, Development and Innovation Fund (HNRDIF) projects GINOP-2.2.1-15-2017-00081, GINOP-2.2.1-15-2017-00033, GINOP-2.3.2-15-2016-00011, EFOP-3.6.2-16-2017-00010, and by the Norway Grant project HU09-0091-A1-2016. RW and GM received support from the projects PD121085 and FK123899 with the support provided from the HNRDIF, financed under the PD16 and FK16 funding schemes. This work was also supported by the János Bolyai Research Scholarship (for GM) of the Hungarian Academy of Sciences.

ACKNOWLEDGMENTS

The support and advice of Professor János Minárovits, Dean Kinga Turzó (Faculty of Dentistry, University of Szeged) are gratefully acknowledged.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.02285/full#supplementary-material>

FIGURE S1 | Rarefaction analysis of the metagenome sequencing of the 10 rumen samples (R1–R10).

FIGURE S2 | Co-occurrence analysis of the metagenome (DNA) samples. Enlargement of the picture shows the taxon belonging to the colored dot in the network. Taxa identified in the core of all ten rumen samples were included in the calculation.

FIGURE S3 | The carbon fixation pathway in prokaryotes (Kegg metabolic map). The blue enzyme numbers have been identified in the RNA dataset.

FIGURE S4 | The propionate metabolism pathway in prokaryotes (Kegg metabolic map). The blue enzyme numbers have been identified in the RNA dataset.

FIGURE S5 | The butanoate metabolism pathway in prokaryotes (Kegg metabolic map). The blue enzyme numbers have been identified in the RNA dataset.

FIGURE S6 | The methanogenesis pathway in prokaryotes (Kegg metabolic map). The blue enzyme numbers have been identified in the RNA dataset.

FIGURE S7 | Co-occurrence analysis of the metatranscriptome (RNA) samples. Enlargement of the picture shows the function belonging to the colored dot in the network. Functions identified in the core of all ten rumen samples were included in the calculation.

TABLE S1 | Composition of feed received by the cows involved in this study.

TABLE S2 | Detailed sequencing data of the metagenomics and metatranscriptomic samples.

TABLE S3 | Compilation of all CAZymes identified in the rumen samples.

TABLE S4 | List of all functions identified in the core, i.e., present in all 10 rumen samples.

REFERENCES

- Al Masalma, M., Raoult, D., and Roux, V. (2009). *Phocaecicola abscessus* gen. nov., sp. nov., an anaerobic bacterium isolated from a human brain abscess sample. *Int. J. Syst. Evol. Microbiol.* 59, 2232–2237. doi: 10.1099/ijs.0.007823-0
- Avagustin, G., Wright, F., and Flint, H. J. (1994). Genetic diversity and phylogenetic relationships among strains of *Prevotella* (*Bacteroides*) *ruminicola* from the rumen. *Int. J. Syst. Bacteriol.* 44, 246–255. doi: 10.1099/00207713-44-2-246
- Bayané, A., and Guio, S. R. (2011). Animal digestive strategies versus anaerobic digestion bioprocesses for biogas production from lignocellulosic biomass. *Rev. Environ. Sci. Biotechnol.* 10, 43–62. doi: 10.1007/s11157-010-9209-4
- Bekele, A. Z., Koike, S., and Kobayashi, Y. (2010). Genetic diversity and diet specificity of ruminal *Prevotella* revealed by 16S rRNA gene-based analysis. *FEMS Microbiol. Lett.* 305, 49–57. doi: 10.1111/j.1574-6968.2010.01911.x
- Berman, K. M., and Chon, M. (1970). Phosphoenolpyruvate synthase of *Escherichia coli*. Purification, some properties, and the role of divalent metal ions. *J. Biol. Chem.* 239, 269–274.
- Bielak, A., Derno, M., Tuchscherer, A., Hammon, H. M., Susenbeth, A., and Kuhla, B. (2016). Body fat mobilization in early lactation influences methane production of dairy cows. *Sci. Rep.* 6:28135. doi: 10.1038/srep28135
- Bognar, A. L., Osborne, C., Shane, B., Singer, S. C., and Ferone, R. (1985). Polyphospho-gamma-glutamate synthetase-dihydrofolate synthetase. Cloning and high expression of the *Escherichia coli* folC gene and purification and properties of the gene product. *J. Biol. Chem.* 260, 5625–5630.
- Bozan, M., Akyol, Ç., Ince, O., Aydin, S., and Ince, B. (2017). Application of next-generation sequencing methods for microbial monitoring of anaerobic digestion of lignocellulosic biomass. *Appl. Microbiol. Biotechnol.* 101, 6849–6864. doi: 10.1007/s00253-017-8438-7
- Brömmelstroet, B. W., Hensgens, C. M. H., Keltjense, J. T., van der Drift, C., and Vogels, D. G. (1990). Purification and properties of 5,10-methylenetetrahydromethanopterin reductase, a Coenzyme F420-dependent enzyme, from *Methanobacterium thermoautotrophicum* strain H. *J. Biol. Chem.* 265, 1652–1657.
- Brömmelstroet, B. W. J., Geerts, W. J., Keltjens, J. T., van der Drift, C., and Vogels, G. D. (1991). Purification and properties of 5,10-methylenetetrahydromethanopterin dehydrogenase and 5,10-methylenetetrahydromethanopterin reductase, two coenzyme F420-dependent enzymes, from *Methanosarcina barkeri*. *Biochim. Biophys. Acta* 1079, 293–302. doi: 10.1016/0167-4838(91)90072-8
- Brul, J. M., Antonopoulos, D. A., Berg Miller, M. E., Wilson, M. K., Yannarell, A. C., Dinsdale, E. A., et al. (2009). Gene-centric metagenomics of the fiber-adherent bovine rumen microbiome reveals forage specific glycoside hydrolases. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1948–1953. doi: 10.1073/pnas.0806191105
- Bryant, M. P., and Burkey, L. A. (1953). Cultural methods and some characteristics of some of the more numerous groups of bacteria in the bovine rumen. *J. Dairy Sci.* 36, 205–217. doi: 10.3168/jds.S0022-0302(53)91482-9
- Bryant, M. P., Small, N., Bouma, C., and Chu, H. (1957). *Bacteroides Ruminicola* n. sp. and *Succinimonas Amylolytica* the new genus and species. *J. Bacteriol.* 76, 15–23.
- Buchfink, B., Xie, C., and Huson, D. H. (2014). Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* 12, 59–60. doi: 10.1038/nmeth.3176
- Campanaro, S., Treu, L., Kougias, P. G., De Francisci, D., Valle, G., and Angelidaki, I. (2016). Metagenomic analysis and functional characterization of the biogas microbiome using high throughput shotgun sequencing and a novel binning strategy. *Biotechnol. Biofuels* 9:26. doi: 10.1186/s13068-016-0441-1
- Case, R. J., Boucher, Y., Dahlöf, L., Holmström, C., Doolittle, W. F., and Kjelleberg, S. (2007). Use of 16S rRNA and rpoB genes as molecular markers for microbial ecology studies. *Appl. Environ. Microbiol.* 73, 278–288. doi: 10.1128/AEM.01177-06
- Chapleur, O., Bize, A., Serain, T., Mazéas, L., and Bouchez, T. (2014). Co-inoculating ruminal content neither provides active hydrolytic microbes nor improves methanization of 13C-cellulose in batch digesters. *FEMS Microbiol. Ecol.* 87, 616–629. doi: 10.1111/1574-6941.12249
- Charon, M. H., Volbeda, A., Chabriere, E., Pieuille, L., and Fontecilla-Camps, J. C. (1999). Structure and electron transfer mechanism of pyruvate:ferredoxin oxidoreductase. *Curr. Opin. Struct. Biol.* 9, 663–669. doi: 10.1016/S0959-440X(99)00027-5
- Chistoserdova, L., Vorholt, J. A., Thauer, R. K., and Lidstrom, M. E. (1998). C1 transfer enzymes and coenzymes linking methylotrophic bacteria and methanogenic Archaea. *Science* 281, 99–102. doi: 10.1126/science.281.5373.99
- Christopherson, M. R., Dawson, J. A., Stevenson, D. M., Cunningham, A. C., Bramhacharya, S., Weimer, P. J., et al. (2014). Unique aspects of fiber degradation by the ruminal ethanologen *Ruminococcus albus* 7 revealed by physiological and transcriptomic analysis. *BMC Genomics* 15:1066. doi: 10.1186/1471-2164-15-1066
- Comtet-Marre, S., Parisot, N., Lepercq, P., Chaucheyras-Durand, F., Mosoni, P., Peyretailade, E., et al. (2017). Metatranscriptomics reveals the active bacterial and eukaryotic fibrolytic communities in the rumen of dairy cow fed a mixed diet. *Front. Microbiol.* 8:67. doi: 10.3389/fmicb.2017.00067
- Cox, M. P., Peterson, D. A., and Biggs, P. J. (2010). SolexaQA: at-a-glance quality assessment of Illumina second-generation sequencing data. *BMC Bioinformatics* 11:485. doi: 10.1186/1471-2105-11-485
- Dai, X., Tian, Y., Li, J., Su, X., Wang, X., Zhao, S., et al. (2015). Metatranscriptomic analyses of plant cell wall polysaccharide degradation by microorganisms in the cow rumen. *Appl. Environ. Microbiol.* 81, 1375–1386. doi: 10.1128/AEM.03682-14
- Deppenmeier, U., Müller, V., and Gottschalk, G. (1996). Pathways of energy conservation in methanogenic Archaea. *Arch. Microbiol.* 165, 149–163. doi: 10.1007/BF01692856
- Deusch, S., Camarinha-Silva, A., Conrad, J., Beifuss, U., Rodehutschord, M., and Seifert, J. (2017). A structural and functional elucidation of the rumen microbiome influenced by various diets and microenvironments. *Front. Microbiol.* 8:1605. doi: 10.3389/fmicb.2017.01605
- Deusch, S., Tilocca, B., Camarinha-Silva, A., and Seifert, J. (2015). News in livestock research - Use of Omics-technologies to study the microbiota in the gastrointestinal tract of farm animals. *Comput. Struct. Biotechnol. J.* 13, 55–63. doi: 10.1016/j.csbj.2014.12.005
- Dijkstra, J. (1994). Production and absorption of volatile fatty acids in the rumen. *Livest. Prod. Sci.* 39, 61–69. doi: 10.1016/0301-6226(94)90154-6
- Dowd, S. E., Callaway, T. R., Wolcott, R. D., Sun, Y., McKeehan, T., Hagevoort, R. G., et al. (2008). Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP). *BMC Microbiol.* 8:125. doi: 10.1186/1471-2180-8-125
- Duncan, S. H., Aminov, R. I., Scott, K. P., Louis, P., Stanton, T. B., and Flint, H. J. (2006). Proposal of *Roseburia faecis* sp. nov., *Roseburia hominis* sp. nov. and *Roseburia inulinivorans* sp. nov., based on isolates from human faeces. *Int. J. Syst. Evol. Microbiol.* 56, 2437–2441. doi: 10.1099/ijs.0.64098-0
- Evans, H. J., and Wood, H. G. (1968). The mechanism of the pyruvate, phosphate dikinase reaction. *Proc. Natl. Acad. Sci. U.S.A.* 61, 1448–1453. doi: 10.1073/pnas.61.4.1448
- Federhen, S. (2015). The NCBI taxonomy database. *Nucleic Acids Res.* 43, D1086–D1098. doi: 10.1093/nar/gku1127
- Ferrer, M., Ghazi, A., Belouqui, A., Vieites, J. M., López-Cortés, N., Marín-Navarro, J., et al. (2012). Functional metagenomics unveils a multifunctional glycosyl hydrolase from the family 43 catalysing the breakdown of plant

- polymers in the calf rumen. *PLoS One* 7:e38134. doi: 10.1371/journal.pone.0038134
- Ferry, J. G. (1999). Enzymology of one-carbon metabolism in methanogenic pathways. *FEMS Microbiol. Lett.* 23, 13–38. doi: 10.1111/j.1574-6976.1999.tb00390.x
- Finn, R. D., Attwood, T. K., Babbitt, P. C., Bateman, A., Bork, P., Bridge, A. J., et al. (2017). InterPro in 2017—beyond protein family and domain annotations. *Nucleic Acids Res.* 45, D190–D199. doi: 10.1093/nar/gkw1107
- Flint, H. J., Bayer, E. A., Rincon, M. T., Lamed, R., and White, B. A. (2008). Polysaccharide utilization by gut bacteria: potential for new insights from genomic analysis. *Nat. Rev. Microbiol.* 6, 121–131. doi: 10.1038/nrmicro1817
- Fondevila, M., and Dehority, B. A. (1996). Interactions between *Fibrobacter succinogenes*, *Prevotella ruminicola*, and *Ruminococcus flavefaciens* in the digestion of cellulose from forages. *J. Anim. Sci.* 74, 678–684. doi: 10.2527/1996.743678x
- Gagen, E. J., Padmanabha, J., Denman, S. E., and McSweeney, C. S. (2015). Hydrogenotrophic culture enrichment reveals rumen *Lachnospiraceae* and *Ruminococcaceae* acetogens and hydrogen-responsive *Bacteroidetes* from pasture-fed cattle. *FEMS Microbiol. Lett.* 362, 1–8. doi: 10.1093/femsle/fnv104
- Godwin, S., Kang, A., Gulino, L.-M., Manefield, M., Gutierrez-Zamora, M.-L., Kienle, M., et al. (2014). Investigation of the microbial metabolism of carbon dioxide and hydrogen in the kangaroo foregut by stable isotope probing. *ISME J.* 8, 1855–1865. doi: 10.1038/ismej.2014.25
- Goodwin, S., McPherson, J. D., and McCombie, W. R. (2016). Coming of age: ten years of next-generation sequencing technologies. *Nat. Rev. Genet.* 17, 333–351. doi: 10.1038/nrg.2016.49
- Güllert, S., Fischer, M. A., Turaev, D., Noebauer, B., Ilmberger, N., Wemheuer, B., et al. (2016). Deep metagenome and metatranscriptome analyses of microbial communities affiliated with an industrial biogas fermenter, a cow rumen, and elephant feces reveal major differences in carbohydrate hydrolysis strategies. *Biotechnol. Biofuels* 9:121. doi: 10.1186/s13068-016-0534-x
- Hackmann, T. J., and Firkins, J. L. (2015). Maximizing efficiency of rumen microbial protein production. *Front. Microbiol.* 6:465. doi: 10.3389/fmicb.2015.00465
- Hallam, S. J., Girguis, P. R., Preston, C. M., Richardson, P. M., and DeLong, E. F. (2003). Identification of methyl coenzyme M reductase A (*mcrA*) genes associated with methane-oxidizing Archaea. *Appl. Environ. Microbiol.* 69, 5483–5491. doi: 10.1128/AEM.69.9.5483-5491.2003
- Hasona, A., Kim, Y., Healy, F. G., Ingram, L. O., and Shanmugam, K. T. (2004). Pyruvate formate lyase and acetate kinase are essential for anaerobic growth of *Escherichia coli* on xylose†. *J. Bacteriol.* 186, 7593–7600. doi: 10.1128/JB.186.22.7593
- Hattori, S. (2008). Syntrophic acetate-oxidizing microbes in methanogenic environments. *Microbes Environ.* 23, 118–127. doi: 10.1264/jsm.2.23.118
- Hegarty, R. S., and Gerdes, R. (1999). Hydrogen production and transfer in the rumen. *Recent Adv. Anim. Nutr. Aust.* 12, 37–44.
- Hellinga, H. W., and Evans, P. R. (1987). Mutations in the active site of *Escherichia coli* phosphofructokinase. *Nature* 327, 437–439. doi: 10.1038/327437a0
- Henderson, G., Cox, F., Ganesh, S., Jonker, A., Young, W., Global Rumen Census Collaborators, et al. (2015). Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci. Rep.* 5:14567. doi: 10.1038/srep14567
- Henderson, G., Leahy, S. C., and Janssen, P. H. (2010). Presence of novel, potentially homoacetogenic bacteria in the rumen as determined by analysis of formyltetrahydrofolate synthetase sequences from ruminants. *Appl. Environ. Microbiol.* 76, 2058–2066. doi: 10.1128/AEM.02580-09
- Hess, M. (2011). Metagenomic discovery of biomass-degrading genes and genomes from cow rumen. *Science* 463, 463–467. doi: 10.1126/science.1200387
- Hobson, P. N. (1969). Rumen bacteria. *Methods Microbiol.* 3, 133–149. doi: 10.1016/S0580-9517(08)70504-X
- Honka, E., Fabry, S., Niemann, T., Palm, P., and Hensel, R. (1990). Properties and primary structure of the L-malate dehydrogenase from the extremely thermophilic archaeobacterium *Methanothermus fervidus*. *Eur. J. Biochem.* 188, 623–632. doi: 10.1111/j.1432-1033.1990.tb15443.x
- Hook, S. E., Wright, A. D. G., and McBride, B. W. (2010). Methanogens: methane producers of the rumen and mitigation strategies. *Archaea* 2010, 50–60. doi: 10.1155/2010/945785
- Hungate, R. E. (1960). Microbial ecology of the rumen. *Bacteriol. Rev.* 24, 353–364. doi: 10.1007/978-94-011-8067-2_9
- Hungate, R. E. (1966). *The Rumen and its Microbes*. New York, NY: Academic Press.
- Huo, W., Zhu, W., and Mao, S. (2014). Impact of subacute ruminal acidosis on the diversity of liquid and solid-associated bacteria in the rumen of goats. *World J. Microbiol. Biotechnol.* 30, 669–680. doi: 10.1007/s11274-013-1489-8
- Huson, D., Auch, A., Qi, J., and Schuster, S. (2007). MEGAN analysis of metagenome data. *Genome Res.* 17, 377–386. doi: 10.1101/gr.5969107
- Huson, D., Mitra, S., and Ruscheweyh, H. (2011). Integrative analysis of environmental sequences using MEGAN4. *Genome Res.* 21, 1552–1560. doi: 10.1101/gr.120618.111.Freely
- Huson, D. H., Beier, S., Flade, I., Górski, A., El-Hadidi, M., Mitra, S., et al. (2016). MEGAN community edition - interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS Comput. Biol.* 12:e1004957. doi: 10.1371/journal.pcbi.1004957
- Imanaka, H., Yamatsu, A., Fukui, T., Atomi, H., and Imanaka, T. (2006). Phosphoenolpyruvate synthase plays an essential role for glycolysis in the modified Embden-Meyerhof pathway in *Thermococcus kodakarensis*. *Mol. Microbiol.* 61, 898–909. doi: 10.1111/j.1365-2958.2006.05287.x
- Jami, E., and Mizrahi, I. (2012). Composition and similarity of bovine rumen microbiota across individual animals. *PLoS One* 7:e33306. doi: 10.1371/journal.pone.0033306
- Janssen, P. H. (2010). Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim. Feed Sci. Technol.* 160, 1–22. doi: 10.1016/j.anifeedsci.2010.07.002
- Janssen, P. H., and Kirs, M. (2008). Structure of the Archaeal community of the rumen. *Appl. Environ. Microbiol.* 74, 3619–3625. doi: 10.1128/AEM.02812-07
- Jewell, K. A., McCormick, C. A., Odt, C. L., Weimer, P. J., and Suen, G. (2015). Ruminal bacterial community composition in dairy cows is dynamic over the course of two lactations and correlates with feed efficiency. *Appl. Environ. Microbiol.* 81, 4697–4710. doi: 10.1128/AEM.00720-15
- Jiang, Y., Xiong, X., Danska, J., and Parkinson, J. (2016). Metatranscriptomic analysis of diverse microbial communities reveals core metabolic pathways and microbiome-specific functionality. *Microbiome* 4:2. doi: 10.1186/s40168-015-0146-x
- Jindou, S., Brulc, J. M., Levy-Assaraf, M., Rincon, M. T., Flint, H. J., Berg, M. E., et al. (2008). Cellulosome gene cluster analysis for gauging the diversity of the ruminal cellulolytic bacterium *Ruminococcus flavefaciens*. *FEMS Microbiol. Lett.* 285, 188–194. doi: 10.1111/j.1574-6968.2008.01234.x
- Jitrapakdee, S., Maurice, M. S., Rayment, I., Cleland, W. W., Wallace, C., and Attwood, P. V. (2008). Structure, mechanism and regulation of pyruvate carboxylase. *Biochem. J.* 413, 369–387. doi: 10.1042/BJ20080709
- Jouany, J. P., and Ushida, K. (1999). The role of protozoa in feed digestion. *Asian Australas. J. Anim. Sci.* 12, 113–128. doi: 10.5713/ajas.1999.113
- Kala, A., Kamra, D. N., Kumar, A., Agarwal, N., Chaudhary, L. C., and Joshi, C. G. (2017). Impact of levels of total digestible nutrients on microbiome, enzyme profile and degradation of feeds in buffalo rumen. *PLoS One* 12:e0172051. doi: 10.1371/journal.pone.0172051
- Kamke, J., Kittelmann, S., Soni, P., Li, Y., Tavendale, M., Ganesh, S., et al. (2016). Rumen metagenome and metatranscriptome analyses of low methane yield sheep reveals a *Sharpea*-enriched microbiome characterised by lactic acid formation and utilisation. *Microbiome* 4:56. doi: 10.1186/s40168-016-0201-2
- Keegan, K. P., Trimble, W. L., Wilkening, J., Wilke, A., Harrison, T., D'Souza, M., et al. (2012). A platform-independent method for detecting errors in metagenomic sequencing data: DRISSEE. *PLoS Comput. Biol.* 8:e1002541. doi: 10.1371/journal.pcbi.1002541
- Kelly, W. J., Henderson, G., Pacheco, D. M., Li, D., Reilly, K., Naylor, G. E., et al. (2016). The complete genome sequence of *Eubacterium limosum* SA11, a metabolically versatile rumen acetogen. *Stand. Genomic Sci.* 11:26. doi: 10.1186/s40793-016-0147-9
- Kim, M., Morrison, M., and Yu, Z. (2011). Status of the phylogenetic diversity census of ruminal microbiomes. *FEMS Microbiol. Ecol.* 76, 49–63. doi: 10.1111/j.1574-6941.2010.01029.x
- King, P. W., Posewitz, M. C., Ghirardi, M. L., and Seibert, M. (2006). Functional studies of [FeFe] hydrogenase maturation in an *Escherichia coli* biosynthetic system. *J. Bacteriol.* 188, 2163–2172. doi: 10.1128/JB.188.6.2163

- Koike, S., and Kobayashi, Y. (2009). Fibrolytic rumen bacteria: their ecology and functions. *Asian Australas. J. Anim. Sci.* 22, 131–138. doi: 10.5713/ajas.2009.r01
- Krause, D. O., Denman, S. E., Mackie, R. I., Morrison, M., Rae, A. L., Attwood, G. T., et al. (2003). Opportunities to improve fiber degradation in the rumen: microbiology, ecology, and genomics. *FEMS Microbiol. Rev.* 27, 663–693. doi: 10.1016/S0168-6445(03)00072-X
- Le Van, T. D., Robinson, J. A., Ralph, J., Greening, R. C., Smolenski, W. J., Leedle, J. A. Z., et al. (1998). Assessment of reductive acetogenesis with indigenous ruminal bacterium populations and *Acetivibrio ruminis*. *Appl. Environ. Microbiol.* 64, 3429–3436.
- Lebioda, L., and Stec, B. (1991). Mechanism of enolase: the crystal structure of enolase-Mg²⁺-2-phosphoglycerate/phosphoenolpyruvate complex at 2.2-Å resolution. *Biochemistry* 30, 2817–2822. doi: 10.1021/bi00225a012
- Lee, C. C., Kibblewhite, R. E., Wagschal, K., Li, R., and Orts, W. J. (2012). Isolation of α -glucuronidase enzyme from a rumen metagenomic library. *Protein J.* 31, 206–211. doi: 10.1007/s10930-012-9391-z
- Li, F., and Guan, L. L. (2017). Metatranscriptomic profiling reveals linkages between the active rumen microbiome and feed efficiency in beef cattle. *Appl. Environ. Microbiol.* 83, 1–16. doi: 10.1128/AEM.00061-17
- Li, R. W., Wu, S., Ransom, L., Baldwin, V. I., Li, W., and Li, C. (2012). Perturbation dynamics of the rumen microbiota in response to exogenous butyrate. *PLoS One* 7:e29392. doi: 10.1371/journal.pone.0029392
- Lin, P. Y., Whang, L. M., Wu, Y. R., Ren, W. J., Hsiao, C. J., Li, S. L., et al. (2007). Biological hydrogen production of the genus *Clostridium*: metabolic study and mathematical model simulation. *Int. J. Hydrogen Energy* 32, 1728–1735. doi: 10.1016/j.ijhydene.2006.12.009
- Liu, C., Meng, Q., Chen, Y., Xu, M., Shen, M., Gao, R., et al. (2017). Role of age-related shifts in rumen bacteria and methanogens in methane production in cattle. *Front. Microbiol.* 8:1563. doi: 10.3389/fmicb.2017.01563
- Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P. M., and Henrissat, B. (2014). The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* 42, 490–495. doi: 10.1093/nar/gkt1178
- Lopez, S., McIntosh, F. M., Wallace, R. J., and Newbold, C. J. (1999). Effect of adding acetogenic bacteria on methane production by mixed rumen microorganisms. *Anim. Feed Sci. Technol.* 78, 1–9. doi: 10.1016/S0377-8401(98)00273-9
- Louis, P., Duncan, S. H., McCrae, S. I., Millar, J., Jackson, M. S., and Flint, H. J. (2004). Restricted distribution of the butyrate kinase pathway among butyrate-producing bacteria from the human colon: restricted distribution of the butyrate kinase pathway among butyrate-producing bacteria from the human colon. *J. Bacteriol.* 186, 2099–2106. doi: 10.1128/JB.186.7.2099
- Lynd, L., Weimer, P., van Zyl, W., and Pretorius, I. (2002). Microbial cellulose utilization: fundamentals and biotechnology. *Microbiol. Mol. Biol. Rev.* 66, 506–577. doi: 10.1128/MMBR.66.3.506
- Lyons, T., Boland, T., Storey, S., and Doyle, E. (2017). Linseed oil supplementation of lambs' diet in early life leads to persistent changes in rumen microbiome structure. *Front. Microbiol.* 8:1656. doi: 10.3389/fmicb.2017.01656
- MacLean, D., Jones, J. D. G., and Studholme, D. J. (2009). Application of “next-generation” sequencing technologies to microbial genetics. *Nat. Rev. Microbiol.* 7, 287–296. doi: 10.1038/nrmicro2088
- Madern, D. (2002). Molecular evolution within the L-malate and L-lactate dehydrogenase super-family. *J. Mol. Evol.* 54, 825–840. doi: 10.1007/s00239-001-0088-8
- Mao, S., Zhang, M., Liu, J., and Zhu, W. (2015). Characterising the bacterial microbiota across the gastrointestinal tracts of dairy cattle: membership and potential function. *Sci. Rep.* 5:16116. doi: 10.1038/srep16116
- Marcus, F., Gontero, B., Harrsch, P. B., and Rittenhouse, J. (1986). Amino acid sequence homology among fructose-1,6-bisphosphatases. *Biochem. Biophys. Res. Commun.* 135, 374–381. doi: 10.1016/0006-291X(86)90005-7
- Martin, C., Morgavi, D. P., and Doreau, M. (2010). Methane mitigation in ruminants: from microbe to the farm scale. *Animal* 4, 351–365. doi: 10.1017/S1751731109990620
- Maspolim, Y., Zhou, Y., Guo, C., Xiao, K., and Ng, W. J. (2015). The effect of pH on solubilization of organic matter and microbial community structures in sludge fermentation. *Bioresour. Technol.* 190, 289–298. doi: 10.1016/j.biortech.2015.04.087
- Matsui, H., Ogata, K., Tajima, K., Nakamura, M., Nagamine, T., Aminov, R. I., et al. (2000). Phenotypic characterization of polysaccharidases produced by four *Prevotella* type strains. *Curr. Microbiol.* 41, 45–49. doi: 10.1007/s002840100089
- Mayorga, O. L., Kingston-Smith, A. H., Kim, E. J., Allison, G. G., Wilkinson, T. J., Hegarty, M. J., et al. (2016). Temporal metagenomic and metabolomic characterization of fresh perennial ryegrass degradation by rumen bacteria. *Front. Microbiol.* 7:1854. doi: 10.3389/fmicb.2016.01854
- Mccann, J. C., Wickersham, T. A., and Loo, J. J. (2014). High-throughput methods redefine the rumen microbiome and its Relationship with Nutrition and Metabolism. *Bioinform. Biol. Insights* 8, 109–125. doi: 10.4137/BBI.S15389
- McDowall, J. S., Murphy, B. J., Haumann, M., Palmer, T., Armstrong, F. A., and Sargent, F. (2014). Bacterial formate hydrogenlyase complex. *Proc. Natl. Acad. Sci. U.S.A.* 111, E3948–E3956. doi: 10.1073/pnas.1407927111
- McSweeney, C. S., Dalrymple, B. P., Gobius, K. S., Kennedy, P. M., Krause, D. O., Mackie, R. I., et al. (1999). The application of rumen biotechnology to improve the nutritive value of fibrous feedstuffs: pre- and post-ingestion. *Livest. Prod. Sci.* 59, 265–283. doi: 10.1016/S0301-6226(99)00032-9
- Meale, S. J., Chaves, A. V., Baah, J., and McAllister, T. A. (2012). Methane production of different forages in in vitro ruminal fermentation. *Asian Australas. J. Anim. Sci.* 25, 86–91. doi: 10.5713/ajas.2011.11249
- Metzker, M. L. (2010). Sequencing technologies - the next generation. *Nat. Rev. Genet.* 11, 31–46. doi: 10.1038/nrg2626
- Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E., Kubal, M., et al. (2011). The metagenomics RAST server: a public resource for the automatic phylogenetic and functional analysis of metagenomes. *Handb. Mol. Microb. Ecol.* 18, 325–331. doi: 10.1002/9781118010518.ch37
- Moran, J. (2005). “How the rumen works,” in *Tropical Dairy Farming Feeding Management Small Holder Dairy Farmers in the Humid Tropics*, (Melbourne: Landlinks Press), 41–49.
- Morgavi, D. P., Forano, E., Martin, C., and Newbold, C. J. (2012). Microbial ecosystem and methanogenesis in ruminants – CORRIGENDUM. *Animal* 6:871. doi: 10.1017/S1751731112000407
- Muirhead, H. (1990). Isoenzymes of pyruvate kinase. *Biochem. Soc. Trans.* 18, 193–196. doi: 10.1042/bst0180193
- Müller, B., Sun, L., Westerholm, M., and Schnürer, A. (2016). Bacterial community composition and fhs profiles of low- and high-ammonia biogas digesters reveal novel syntrophic acetate-oxidising bacteria. *Biotechnol. Biofuels* 9:48. doi: 10.1186/s13068-016-0454-9
- Naas, A. E., Mackenzie, A. K., Mravec, J., Schückel, J., Willats, W. G. T., Eijssink, V. G. H., et al. (2014). Do rumen Bacteroidetes utilize an alternative mechanism for cellulose degradation? *mBio* 5:e01401-14. doi: 10.1128/mBio.01401-14
- Nagai, F., Watanabe, Y., and Morotomi, M. (2010). *Slackia piriformis* sp. nov. and *Collinsella tanakaei* sp. nov., new members of the family *Coriobacteriaceae*, isolated from human faeces. *Int. J. Syst. Evol. Microbiol.* 60, 2639–2646. doi: 10.1099/ijs.0.017533-0
- Nakayama, T., Yonekura, S.-I., Yonei, S., and Zhang-Akiyama, Q.-M. (2013). *Escherichia coli* pyruvate:flavodoxin oxidoreductase, YdbK - regulation of expression and biological roles in protection against oxidative stress. *Genes Genet. Syst.* 88, 175–188. doi: 10.1266/ggs.88.175
- Opio, C., Gerber, P., Mottet, A., Falcucci, A., Tempio, G., MacLeod, M., et al. (2013). *Greenhouse gas Emissions from Ruminant Supply Chains—A Global Life Cycle Assessment*. Available at: <http://www.fao.org/docrep/018/i3461e/i3461e.pdf>
- Oulas, A., Pavloudi, C., Polymenakou, P., Pavlopoulos, G. A., Papanikolaou, N., Kotoulas, G., et al. (2015). Metagenomics: tools and insights for analyzing next-generation sequencing data derived from biodiversity studies. *Bioinform. Biol. Insights* 9, 75–88. doi: 10.4137/BBI.S12462
- Parmar, N. R., Nirmal Kumar, J. I., and Joshi, C. G. (2015). Deep insights into carbohydrate metabolism in the rumen of Mehsani buffalo at different diet treatments. *Genomics Data* 6, 59–62. doi: 10.1016/j.gdata.2015.08.007
- Patel, D. D., Patel, A. K., Parmar, N. R., Shah, T. M., Patel, J. B., Pandya, P. R., et al. (2014). Microbial and Carbohydrate Active Enzyme profile of buffalo rumen metagenome and their alteration in response to variation in the diet. *Gene* 545, 88–94. doi: 10.1016/j.gene.2014.05.003
- Petri, R. M., Schwaiger, T., Penner, G. B., Beauchemin, K. A., Forster, R. J., McKinnon, J. J., et al. (2013). Characterization of the core rumen microbiome in cattle during transition from forage to concentrate as well as during and after an acidotic challenge. *PLoS One* 8:e83424. doi: 10.1371/journal.pone.0083424

- Picard, C., Fioramonti, J., Francois, A., Robinson, T., Neant, F., and Matuchansky, C. (2005). Review article: bifidobacteria as probiotic agents - Physiological effects and clinical benefits. *Aliment. Pharmacol. Ther.* 22, 495–512. doi: 10.1111/j.1365-2036.2005.02615.x
- Pitta, D. W., Indugu, N., Kumar, S., Vecchiarelli, B., Sinha, R., Baker, L. D., et al. (2016). Metagenomic assessment of the functional potential of the rumen microbiome in Holstein dairy cows. *Anaerobe* 38, 50–60. doi: 10.1016/j.anaerobe.2015.12.003
- Pitta, D. W., Parmar, N., Patel, A. K., Indugu, N., Kumar, S., Prajapati, K. B., et al. (2014). Bacterial diversity dynamics associated with different diets and different primer pairs in the rumen of Kankrej cattle. *PLoS One* 9:e111710. doi: 10.1371/journal.pone.0111710
- Pomper, B. K., Vorholt, J. A., Chistoserdova, L., Lidstrom, M. E., and Thauer, R. K. (1999). A methenyl tetrahydromethanopterin cyclohydrolase and a methenyl tetrahydrofolate cyclohydrolase in *Methylobacterium extorquens* AM1. *Eur. J. Biochem.* 261, 475–480. doi: 10.1046/j.1432-1327.1999.00291.x
- Pope, P. B., Mackenzie, A. K., Gregor, I., Smith, W., Sundset, M. A., McHardy, A. C., et al. (2012). Metagenomics of the svalbard reindeer rumen microbiome reveals abundance of polysaccharide utilization loci. *PLoS One* 7:e38571. doi: 10.1371/journal.pone.0038571
- Poulsen, M., Schwab, C., Borg Jensen, B., Engberg, R. M., Spang, A., Canibe, N., et al. (2013). Methylophilic methanogenic Thermoplasmata implicated in reduced methane emissions from bovine rumen. *Nat. Commun.* 4:1428. doi: 10.1038/ncomms2432
- Purush, J., Fouts, D. E., Morrison, M., White, B. A., Mackie, R. I., Coutinho, P. M., et al. (2010). Comparative genome analysis of *Prevotella ruminicola* and *Prevotella bryantii*: insights into their environmental niche. *Microb. Ecol.* 60, 721–729. doi: 10.1007/s00248-010-9692-8
- Qiao, G. H., Shan, A. S., Ma, N., Ma, Q. Q., and Sun, Z. W. (2010). Effect of supplemental *Bacillus* cultures on rumen fermentation and milk yield in Chinese Holstein cows. *J. Anim. Physiol. Anim. Nutr.* 94, 429–436. doi: 10.1111/j.1439-0396.2009.00926.x
- Raben, N., Exelbert, R., Spiegel, R., Sherman, J. B., Nakajima, H., Plotz, P., et al. (1995). Functional expression of human mutant phosphofructokinase in yeast: genetic defects in French Canadian and Swiss patients with phosphofructokinase deficiency. *Am. J. Hum. Genet.* 56, 131–141.
- Raes, J., Foerster, K. U., and Bork, P. (2007). Get the most out of your metagenome: computational analysis of environmental sequence data. *Curr. Opin. Microbiol.* 10, 490–498. doi: 10.1016/j.mib.2007.09.001
- Ragsdale, S. W., and Pierce, E. (2009). Acetogenesis and the Wood-Ljungdahl pathway of CO₂ fixation. *Biochim. Biophys. Acta* 1784, 1873–1898. doi: 10.1016/j.bbapap.2008.08.012
- Rajendran, J., and Gunasekaran, P. (2011). Microbial phylogeny and diversity: small subunit ribosomal RNA sequence analysis and beyond. *Microbiol. Res.* 166, 99–110. doi: 10.1016/j.micres.2010.02.003
- Ramšak, A., Peterka, M., Tajima, K., Martin, J. C., Wood, J., Johnston, M. E. A., et al. (2000). Unravelling the genetic diversity of ruminal bacteria belonging to the CFB phylum. *FEMS Microbiol. Ecol.* 33, 69–79. doi: 10.1016/S0168-6496(00)00049-0
- Ravachol, J., Borne, R., Meynial-Salles, I., Soucaille, P., Pagès, S., Tardif, C., et al. (2015). Combining free and aggregated cellulolytic systems in the cellulosome-producing bacterium *Ruminiclostridium cellulolyticum*. *Biotechnol. Biofuels* 8:114. doi: 10.1186/s13068-015-0301-4
- Reichardt, N., Duncan, S. H., Young, P., Belenguer, A., McWilliam Leitch, C., Scott, K. P., et al. (2014). Phylogenetic distribution of three pathways for propionate production within the human gut microbiota. *ISME J.* 8, 1323–1335. doi: 10.1038/ismej.2014.14
- Rock, C. O., and Jackowski, S. (2002). Forty years of bacterial fatty acid synthesis. *Biochem. Biophys. Res. Commun.* 292, 1155–1166. doi: 10.1006/bbrc.2001.2022
- Rosewarne, C. P., Pope, P. B., Cheung, J. L., and Morrison, M. (2014). Analysis of the bovine rumen microbiome reveals a diversity of Sus-like polysaccharide utilization loci from the bacterial phylum Bacteroidetes. *J. Ind. Microbiol. Biotechnol.* 41, 601–606. doi: 10.1007/s10295-013-1395-y
- Russell, J. B., and Rychlik, J. L. (2008). Factors that alter rumen microbial ecology. *Science* 292, 1119–1122. doi: 10.1126/science.1058830
- Samuelov, N. S., Lamed, R., Lowe, S., and Zeikus, J. G. (1991). Influence of CO₂-HCO₃⁻ levels and pH on growth, succinate production, and enzyme activities of *Anaerobiospirillum succiniciproducens*. *Appl. Environ. Microbiol.* 57, 3013–3019. doi: 10.1007/s00253-014-6014-y
- Schären, M., Kiri, K., Riede, S., Gardener, M., Meyer, U., Hummel, J., et al. (2017). Alterations in the rumen liquid-, particle- and epithelium-associated microbiota of dairy cows during the transition from a silage- and concentrate-based ration to pasture in spring. *Front. Microbiol.* 8:744. doi: 10.3389/fmicb.2017.00744
- Schöcke, L., and Weimer, P. J. (1997). Purification and characterization of phosphoenolpyruvate carboxykinase from the anaerobic ruminal bacterium *Ruminococcus flavefaciens*. *Arch. Microbiol.* 167, 289–294. doi: 10.1007/s002030050446
- Schuchmann, K., and Müller, V. (2014). Autotrophy at the thermodynamic limit of life: a model for energy conservation in acetogenic bacteria. *Nat. Rev. Microbiol.* 12, 809–821. doi: 10.1038/nrmicro3365
- Seo, J. K., Park, T. S., Kwon, I. H., Piao, M. Y., Lee, C. H., and Ha, J. K. (2013). Characterization of cellulolytic and xylanolytic enzymes of *Bacillus licheniformis* JK7 isolated from the rumen of a native Korean goat. *Asian Australas. J. Anim. Sci.* 26, 50–58. doi: 10.5713/ajas.2012.12506
- Shabat, S. K., Sasson, G., Doron-Faigenboim, A., Durman, T., Yaacoby, S., Berg Miller, M. E., et al. (2016). Specific microbiome-dependent mechanisms underlie the energy harvest efficiency of ruminants. *ISME J.* 10, 1–15. doi: 10.1038/ismej.2016.62
- Shen, H., Lu, Z., Xu, Z., Chen, Z., and Shen, Z. (2017). Associations among dietary non-fiber carbohydrate, ruminal microbiota and epithelium G-protein-coupled receptor, and histone deacetylase regulations in goats. *Microbiome* 5:123. doi: 10.1186/s40168-017-0341-z
- Shi, W., Moon, C. D., Leahy, S. C., Kang, D., Froula, J., Kittelmann, S., et al. (2014). Methane yield phenotypes linked to differential gene expression in the sheep rumen microbiome. *Genome Res.* 24, 1517–1525. doi: 10.1101/gr.168245.113
- Shinkai, T., Mitsumori, M., Sofyan, A., Kanamori, H., Sasaki, H., Katayose, Y., et al. (2016). Comprehensive detection of bacterial carbohydrate-active enzyme coding genes expressed in cow rumen. *Anim. Sci. J.* 87, 1363–1370. doi: 10.1111/asj.12585
- Singh, K. M., Reddy, B., Patel, A. K., Panchasara, H., Parmar, N., Patel, A. B., et al. (2014). Metagenomic analysis of buffalo rumen microbiome: effect of roughage diet on Dormancy and Sporulation genes. *Meta Gene* 2, 252–268. doi: 10.1016/j.mgene.2014.01.005
- Siriwongrungron, V., Zeng, R. J., and Angelidaki, I. (2007). Homoacetogenesis as the alternative pathway for H₂ sink during thermophilic anaerobic degradation of butyrate under suppressed methanogenesis. *Water Res.* 41, 4204–4210. doi: 10.1016/j.watres.2007.05.037
- Skillman, L. C., Skillman, L. C., Toovey, A. F., Toovey, A. F., Williams, A. J., and Williams, A. J. (2006). Development and validation of a real-time PCR method to quantify rumen protozoa and examination of variability between. *Cloning* 72, 200–206. doi: 10.1128/AEM.72.1.200
- Stevens, C. E., and Hume, I. D. (1998). Contributions of microbes in vertebrate gastrointestinal tract to production and conservation of nutrients. *Physiol. Rev.* 78, 393–427. doi: 10.1152/physrev.1998.78.2.393
- Stevenson, D. M., and Weimer, P. J. (2007). Dominance of *Prevotella* and low abundance of classical ruminal bacterial species in the bovine rumen revealed by relative quantification real-time PCR. *Appl. Microbiol. Biotechnol.* 75, 165–174. doi: 10.1007/s00253-006-0802-y
- Stewart, R. D., Auffret, M. D., Warr, A., Wiser, A. H., Press, M. O., Langford, K. W., et al. (2018). Assembly of 913 microbial genomes from metagenomic sequencing of the cow rumen. *Nat. Commun.* 9:870. doi: 10.1038/s41467-018-03317-6
- Svartström, O., Alneberg, J., Terrapon, N., Lombard, V., De Bruijn, I., Malmsten, J., et al. (2017). Ninety-nine de novo assembled genomes from the moose (*Alces alces*) rumen microbiome provide new insights into microbial plant biomass degradation. *ISME J.* 11, 2538–2551. doi: 10.1038/ismej.2017.108
- Takahashi-Iniguez, T., García-Hernandez, E., Arreguín-Espinosa, R., and Flores, M. E. (2012). Role of vitamin B12 on methylmalonyl-CoA mutase activity. *J. Zhejiang Univ. Sci. B* 13, 423–437. doi: 10.1631/jzus.B1100329
- Thoetkiattikul, H., Mhuantong, W., Laothanachareon, T., Tangphatsornruang, S., Pattarajinda, V., Eurwilaichitr, L., et al. (2013). Comparative analysis of microbial profiles in cow rumen fed with different dietary fiber by tagged 16S rRNA gene pyrosequencing. *Curr. Microbiol.* 67, 130–137. doi: 10.1007/s00284-013-0336-3

- Tyson, G. W., Chapman, J., Hugenholtz, P., Allen, E. E., Ram, R. J., Richardson, P. M., et al. (2004). Community structure and metabolism through reconstruction of microbial genomes from the environment. *Nature* 428, 37–43. doi: 10.1038/nature02340
- Ungerfeld, E. M. (2014). A theoretical comparison between two ruminal electron sinks. *Front. Microbiol.* 5:235. doi: 10.3389/fmicb.2014.00235
- Ungerfeld, E. M. (2015). Shifts in metabolic hydrogen sinks in the methanogenesis-inhibited ruminal fermentation: a meta-analysis. *Front. Microbiol.* 6:37. doi: 10.3389/fmicb.2015.00037
- Ushakova, N. A., Nekrasov, R. V., Meleshko, N. A., Laptev, G. Y., Il'ina, L. A., Kozlova, A. A., et al. (2013). Effect of *Bacillus subtilis* on the rumen microbial community and its components exhibiting high correlation coefficients with the host nutrition, growth, and development. *Microbiology* 82, 475–481. doi: 10.1134/S0026261713040127
- Venter, J. C., Adams, M. D., Myers, E. W., Li, P. W., Mural, R. J., Sutton, G. G., et al. (2001). The sequence of the human genome. *Science* 291, 1304–1351. doi: 10.1126/science.1058040
- Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., et al. (2004). Environmental genome shotgun sequencing of the Sargasso sea. *Science* 304, 66–74. doi: 10.1126/science.1093857
- Wallace, R. J., Rooke, J. A., McKain, N., Duthie, C.-A., Hyslop, J. J., Ross, D. W., et al. (2015). The rumen microbial metagenome associated with high methane production in cattle. *BMC Genomics* 16:839. doi: 10.1186/s12864-015-2032-0
- Wallace, R. J., Snelling, T. J., McCartney, C. A., Tapio, I., and Strozzi, F. (2017). Application of meta-omics techniques to understand greenhouse gas emissions originating from ruminal metabolism. *Genet. Sel. Evol.* 49, 1–11. doi: 10.1186/s12711-017-0285-6
- Wang, L., Hatem, A., Catalyurek, U. V., Morrison, M., and Yu, Z. (2013). Metagenomic insights into the carbohydrate-active enzymes carried by the microorganisms adhering to solid digesta in the rumen of cows. *PLoS One* 8:e78507. doi: 10.1371/journal.pone.0078507
- Wang, W., Li, C., Li, F., Wang, X., Zhang, X., Liu, T., et al. (2016). Effects of early feeding on the host rumen transcriptome and bacterial diversity in lambs. *Sci. Rep.* 6:32479. doi: 10.1038/srep32479
- Wegener, G., and Krause, U. (2002). Different modes of activating phosphofructokinase, a key regulatory enzyme of glycolysis, in working vertebrate muscle. *Biochem. Soc. Trans.* 30, 264–270. doi: 10.1042/BST0300264
- Weimer, P. J. (2015). Redundancy, resilience, and host specificity of the ruminal microbiota: implications for engineering improved ruminal fermentations. *Front. Microbiol.* 6:296. doi: 10.3389/fmicb.2015.00296
- Wen, Z., Morrison, M., and Wen, Z. (1996). Characterization of ammonia utilization by *Prevotella ruminicola* B 14. *Anim. Sci.* 496, 1–3.
- Williams, L., Nguyen, T., Li, Y., Porter, T. N., and Raushel, F. M. (2006). Uronate isomerase: a nonhydrolytic member of the amidohydrolase superfamily with an ambivalent requirement for a divalent metal ion. *Biochemistry* 45, 7453–7462. doi: 10.1007/s11103-011-9767-z
- Wirth, R., Kovács, E., Maróti, G., Bagi, Z., Rákhely, G., and Kovács, K. L. (2012). Characterization of a biogas-producing microbial community by short-read next generation DNA sequencing. *Biotechnol. Biofuels* 5:41. doi: 10.1186/1754-6834-5-41
- Wirth, R., Lakatos, G., Böjti, T., Maróti, G., Bagi, Z., Kis, M., et al. (2015a). Metagenome changes in the mesophilic biogas-producing community during fermentation of the green alga *Scenedesmus obliquus*. *J. Biotechnol.* 215, 52–61. doi: 10.1016/j.jbiotec.2015.06.396
- Wirth, R., Lakatos, G., Maróti, G., Bagi, Z., Minárovics, J., Nagy, K., et al. (2015b). Exploitation of algal-bacterial associations in a two-stage biohydrogen and biogas generation process. *Biotechnol. Biofuels* 8:59. doi: 10.1186/s13068-015-0243-x
- Woese, C. R., and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. U.S.A.* 74, 5088–5090. doi: 10.1073/pnas.74.11.5088
- Wu, S., Baldwin, R. L., Li, W., Li, C., Connor, E. E., and Li, R. W. (2012). The bacterial community composition of the bovine rumen detected using pyrosequencing of 16S rRNA genes. *Metagenomics* 1, 1–11. doi: 10.4303/mg/235571
- Xu, K., Liu, H., Du, G., and Chen, J. (2009). Real-time PCR assays targeting formyltetrahydrofolate synthetase gene to enumerate acetogens in natural and engineered environments. *Anaerobe* 15, 204–213. doi: 10.1016/j.anaerobe.2009.03.005
- Yamada, Y., and Yukphan, P. (2008). Genera and species in acetic acid bacteria. *Int. J. Food Microbiol.* 125, 15–24. doi: 10.1016/j.jfoodmicro.2007.11.077
- Yuan, J., Doucette, C. D., Fowler, W. U., Feng, X.-J., Piazza, M., Rabitz, H. A., et al. (2009). Metabolomics-driven quantitative analysis of ammonia assimilation in *E. coli*. *Mol. Syst. Biol.* 5:302. doi: 10.1038/msb.2009.60
- Zhang, J., Shi, H., Wang, Y., Li, S., Cao, Z., Ji, S., et al. (2017). Effect of dietary forage to concentrate ratios on dynamic profile changes and interactions of ruminal microbiota and metabolites in holstein heifers. *Front. Microbiol.* 8:2206. doi: 10.3389/fmicb.2017.02206

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Wirth, Kádár, Kakuk, Maróti, Bagi, Szilágyi, Rákhely, Horváth and Kovács. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.